



EVIDENCE FOR ACTION TECHNICAL PAPERS

INTERVENTIONS TO ADDRESS HIV IN PRISONS NEEDLE AND SYRINGE PROGRAMMES AND DECONTAMINATION STRATEGIES



World Health
Organization



UNITED NATIONS
Office on Drugs and Crime



UNAIDS
JOINT UNITED NATIONS PROGRAMME ON HIV/AIDS

UNHCR
UNICEF
WFP
UNDP
UNFPA
UNODC
ILO
UNESCO
WHO
WORLD BANK

WHO Library Cataloguing-in-Publication Data

Interventions to address HIV in prisons: needle and syringe programmes and decontamination strategies /
Ralf Jürgens.

(Evidence for Action Technical Papers)

1.HIV infections – prevention and control. 2.Acquired immunodeficiency syndrome – prevention and control. 3.Substance Abuse, Intravenous – complications. 4.Needle exchange programs. 5.Prisons. I.World Health Organization. II.UNAIDS. III.UNODC. IV.Jürgens, Ralf.

ISBN 978 92 4 159581 0

(NLM classification: WC 503.6)

© **World Health Organization 2007**

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; e-mail: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

The named author alone is responsible for the views expressed in this publication.

Printed in Geneva

EVIDENCE FOR ACTION TECHNICAL PAPERS

INTERVENTIONS TO ADDRESS HIV IN PRISONS

NEEDLE AND SYRINGE PROGRAMMES AND DECONTAMINATION STRATEGIES

World Health Organization, UNODC, UNAIDS / Geneva, 2007

ACKNOWLEDGEMENTS

This document and the other components of the Evidence for Action Technical Paper on Effectiveness of Interventions to Address HIV in Prisons were authored by Ralf Jürgens.

The review of the evidence was commissioned by Sandra Black (WHO Department of HIV/AIDS). A large international network of researchers and colleagues made important contributions to the development of these publications. The accompanying references illustrate the seminal contributions of many investigators and their colleagues who have contributed to the science and practice of HIV interventions in prisons. Particular thanks go to Kate Dolan and colleagues who have undertaken much groundbreaking research and published extensively in this area; Anke Stallwitz and Heino Stöver who have recently completed a literature review on substitution treatment in prisons; Emanuele Pontali for his excellent review of issues related to provision of antiretroviral therapy in prison; Rick Lines for his work on prison-based needle and syringe programmes; and Judy Auerbach, Monica Beg, Dave Burrows, Holly Catania, Paddy Costall, Anindya Chatterjee, Martin Donoghoe, Joumana Hermez, Jothi Raja, Christian Kroll, Morag MacDonald, Martina Melis, Lars Moller, Igor Oliynyk, Jyothi Raja, Gray Sattler, Bobby Smyth, Gerald Thomas, Mike Trace, Ron Valdiserri, Alex Wodak, and the Community Acquired Infections Division of the Public Health Agency of Canada for comments provided on (parts of) a first draft of the publications.

We acknowledge the many individuals from many countries who have attempted to fund and encourage research in this area, assist the difficult process of translating research findings into policy, operate services despite funding inadequacy and work with prisoners and the communities they come from and to whom most of them return; and the cooperation of prisoners who, over the years, have participated in the studies and interventions and have taught the public health community about their lives in prison and how to effectively manage HIV in prisons.

Annette Verster and Andrew Ball at WHO Department of HIV/AIDS, and Andrew Doupe, HIV and Legal Consultant, edited the publication, which was developed under the supervision of Jos Perriëns, WHO Department of HIV/AIDS.

WHO wishes to acknowledge the generous contributions of the Australian Agency for International Development and the Dutch Ministry of Health to the development of this document.

CONTENTS

Preface	4
Executive summary	5
Methodology	7
1. Evidence regarding injecting drug use and resulting transmission of HIV and other blood-borne infections in prisons.....	8
1.1 Background	8
1.2 Injecting drug use in prison	8
1.2.1 Starting to inject in prison.....	9
1.2.2 Using non-sterile injecting equipment in prison.....	9
1.2.3 Determinants of injecting drug use in prison.....	10
1.2.4 Injecting upon release.....	11
1.3 Evidence of HIV and HCV transmission resulting from injecting in prison	11
2. Evidence regarding prison-based needle and syringe programmes.....	12
2.1 Background	12
2.2 Evidence of effectiveness of NSPs in community settings	12
2.3 Evidence of the effectiveness of NSPs in prison.....	13
2.3.1 Reduction in the use of non-sterile injecting equipment and resulting blood-borne infections.....	13
2.3.2 Additional benefits.....	13
2.3.3 Absence of unintended negative consequences	14
2.3.4 Other findings	15
2.4 Conclusions and recommendations.....	16
3. Evidence regarding bleach and decontamination strategies.....	19
3.1 Background	19
3.2 Evidence of effectiveness of bleach programmes in the community	19
3.3 Evidence from studies undertaken in prisons	19
3.3.1 Reduction of risk behaviours and of infections.....	19
3.3.2 Safety and security.....	20
3.4 Conclusions and Recommendations.....	20
Table 1: Examples of studies that have examined injecting behaviour in prison.....	21
Table 2: Association of HIV, HCV and HBV among injecting drug users with a history of imprisonment.....	22
Table 3: Countries with needle and syringe programmes in prisons	25
Table 4: Sample evaluations of needle and syringe programmes in prisons.....	25
References	26

PREFACE

The global environment for the HIV response has shifted substantially towards a massive scaling up of prevention, treatment and care interventions. In particular, Governments made an unprecedented commitment during the United Nations Special Session on HIV/AIDS in 2001 to halting and reversing the epidemic by 2015. More recently, at the 2005 World Summit and at the 2006 High Level Meeting on AIDS, Governments committed to pursue all necessary efforts towards the goal of universal access to comprehensive prevention programmes, treatment, care and support by 2010. In support of this, substantial additional resources to fund an expanded response have become available, including through the Global Fund to Fight AIDS, Tuberculosis and Malaria.

Governments face the challenge of translating these commitments into practical programmes, which includes implementing a comprehensive range of interventions to address HIV transmission related to injecting drug use, including in their prison systems. This publication is part of a series of Evidence for Action Technical Papers, which aim to make the evidence for the effectiveness of interventions to manage HIV in prisons accessible to policy-makers and programmers. The series consists of:

1. Four papers that consider the effectiveness of a number of key interventions in managing HIV in

prisons, including:

- needle and syringe programmes and decontamination strategies;
- prevention of sexual transmission;
- drug dependence treatments; and
- HIV care, treatment and support.

2. A comprehensive paper on *Effectiveness of Interventions to Address HIV in Prisons* which (1) provides much more detailed information about the interventions covered in the four above mentioned papers; and (2) reviews the evidence regarding HIV prevalence, risk behaviours and transmission in prisons, as well as other interventions that are part of a comprehensive approach to managing HIV in prisons, including HIV education, testing and counselling, and other programmes. This paper is available, in electronic format only, at <http://www.who.int/hiv/idu/en>.

WHO, UNODC and UNAIDS recognize the importance of this review in supporting the implementation and scale up of evidence-based interventions in prison settings aimed at HIV prevention, treatment and care.

A NOTE ON TERMINOLOGY

In some jurisdictions different terms are used to denote places of detention, which hold people who are awaiting trial, who have been convicted or who are subject to other conditions of security. Similarly, different words are used for various groups of people who are detained.

In this paper, the term '**prison**' has been used for all places of detention and the term '**prisoner**' has been used to describe all who are held in such places, including adult and juvenile males and females detained in criminal justice and prison facilities during the investigation of a crime; while awaiting trial; after conviction and before sentencing; and after sentencing. Although the term does not formally cover persons detained for reasons relating to **immigration or refugee status**, those **detained without charge**, and those sentenced to **compulsory treatment and rehabilitation centres** as they exist in some countries, nonetheless most of the considerations in this paper apply to them as well.

In this paper, the term '**needle and syringe programmes**' (**NSPs**) refers to programmes that provide people who inject drugs with access to sterile injecting equipment (needles and syringes, swabs, vials of sterile water) and most often also to health education, referrals, counselling and other services. This term has grown in popularity and is increasingly replacing terms such as "needle exchange programmes" or "syringe exchange programmes." In prisons, in some NSPs used injecting equipment is exchanged for new injecting equipment, for example through automated machines. However, in most prison-based NSPs, as in the community, injecting equipment is distributed, information about and the means for the safe disposal of syringes are provided, and additional services are also offered.

EXECUTIVE SUMMARY

HIV hit prisons early and hit them hard. The rates of HIV infection among prisoners in many countries are significantly higher than those in the general population. Hepatitis C virus (HCV) seroprevalence rates are even higher. While most of the prisoners living with HIV or AIDS in prison contract their infection outside the institutions before imprisonment, the risk of being infected in prison, in particular through sharing of contaminated injecting equipment and through unprotected sex, is high. Studies from around the world show that injecting drug use is a reality in many prisons and that most prisoners who inject have to share injecting equipment, creating a serious risk of spread of infection. Even countries that have invested heavily in drug demand and drug supply reduction efforts in prisons have not been able to stop injecting drug use. Outbreaks of HIV infection have occurred in a number of prison systems, demonstrating how rapidly HIV can spread in prison unless effective action is taken to prevent transmission.

The importance of implementing HIV interventions, including needle and syringe programmes, in prisons was recognized early in the epidemic. After holding a first consultation on prevention and control of HIV in prisons in 1987, WHO responded to growing evidence of HIV infection in prisons worldwide by issuing guidelines on HIV infection and AIDS in prisons in 1993. With regard to health care and prevention of HIV, the guidelines emphasize that “all prisoners have the right to receive health care, including preventive measures, equivalent to that available in the community without discrimination, in particular with respect to their legal status or nationality”. In particular, the guidelines recommend that “in countries where clean syringes and needles are made available to injecting drug users in the community, consideration should be given to providing clean injection equipment during detention and on release”. Such recommendations were recently re-affirmed in the 2006 framework for an effective national response to HIV/AIDS in prisons, jointly published by the United Nations Office on Drugs and Crime (UNODC), WHO, and UNAIDS.

An increasing number of countries has introduced HIV programmes in prisons since the early 1990s. However, many of them are small in scale, restricted to a few prisons, or exclude necessary interventions for which evidence of effectiveness exists. There is an urgent need to introduce comprehensive programmes (including information and education, particularly through peers; drug dependence treatment, in particular opioid substitution therapy with metha-

done and/or buprenorphine; provision of condoms; diagnosis and treatment of STIs, voluntary counselling and HIV testing; and HIV care and support, including provision of antiretroviral treatment), and to scale them up rapidly. As part of these programmes, prison systems should consider introducing needle and syringe programmes.

Needle and syringe programmes

There is evidence that **needle and syringe programmes** (NSPs) are **feasible** in a wide range of prison settings, including in men’s and women’s prisons, prisons of all security levels, and small and large prisons. There is evidence that providing clean needles and syringes is readily accepted by IDUs in prisons and that it contributes to a significant reduction of syringe sharing over time. It also appears to be effective in reducing resulting HIV infections. At the same time, there is no evidence to suggest that prison-based NSPs have serious, unintended negative consequences. In particular, they do not appear to lead to increased drug use or injecting, nor are they used as weapons. Evaluations have found that NSPs in prisons actually facilitate referral of drug users to drug dependence treatment programmes. Ultimately, since most prisoners leave prison at some point to return to their community, implementing NSPs in prisons will benefit not only prisoners and prison staff, but also society in general. Therefore, it is recommended that

- ▶ **Prison authorities in countries experiencing or threatened by an epidemic of HIV infections among IDUs should introduce NSPs urgently and expand implementation to scale as soon as possible.** The higher the prevalence of injecting drug use and associated risk behaviour is in prison, the more urgent introduction of prison-based NSPs becomes.
- ▶ **Prisoners should have easy, confidential access to NSPs**, and prisoners and staff should receive information and education about the programmes and be involved in their design and implementation.
- ▶ **Carefully evaluated pilot programmes of prison-based NSPs may be important in allowing the introduction of these programmes, but they should not delay the expansion of the programmes**, particularly where there already is evidence of high levels of injecting in prisons.
- ▶ **Additional research about prison-based NSPs should be undertaken.** In particular,

more research in resource-poor systems outside Western Europe could allow for more rapid expansion of NSPs in these settings. Research should be designed to address operational issues and research gaps rather than replicate existing studies. Evaluation of pilot programmes may be justified if: (1) the evaluation takes place in settings that are sufficiently different from settings in which evaluations have already been undertaken; or (2) it addresses research gaps.

Bleach and decontamination strategies

Evaluations of **bleach** programmes in prisons have shown that distribution of bleach or other disinfectants is feasible in prisons and does not compromise security. However, disinfection and decontamination schemes in the community outside prisons **are not supported by evidence of effectiveness**. Studies undertaken in prisons have shown that conditions in prisons further reduce the probability that injecting equipment may be effectively decontaminated. **Because of their limited effectiveness, bleach programmes can only be regarded as a second-line strategy to NSPs.** Therefore:

- ▶ Bleach programmes should be available in prisons where authorities continue to oppose the introduction of NSPs despite evidence of their effectiveness, and to complement NSPs. However, they cannot replace NSPs.
- ▶ Where bleach programmes are implemented, bleach should be made easily and discreetly accessible to prisoners in various locations in the prison, together with information and education about how to clean injecting equipment and information about the limited efficacy of bleach as a disinfectant for inactivating HIV and particularly HCV.
- ▶ Where bleach programmes exist in prisons, but not NSPs, public health practitioners should continue to advocate for the introduction of NSPs.

METHODOLOGY

A comprehensive search of the published literature was carried out. Electronic library and HIV/AIDS databases, and websites of various government and non-governmental bodies, relevant conferences, and prison health and health news sites were searched. Key search terms used included “prison(s)”, “jail(s)”, “detention centre(s)”, “correctional facility(ies)”, “prisoner(s)”, inmate(s), “HIV”, “human immunodeficiency virus”, “hepatitis C”, and “HCV”. These search terms were combined with specific interventions (such as “condom(s)”, “bleach”, “needle exchange” etc) and, where useful, with specific countries or regions. Studies and other materials reported in English, French, German, Italian, Portuguese and Spanish were reviewed. Attempts were made to access information from developing countries and to access the ‘grey’ literature through professional contacts, and direct contact with known researchers and research centres. Nevertheless, the review had limitations: not all papers could be obtained and publications in languages other than those mentioned are not included.

Generally, the review examines whether interventions to manage HIV in prisons have been demonstrated scientifically to reduce the spread of HIV among prisoners or to have other positive health effects. The evidence has been evaluated according to the criteria originally proposed by Bradford Hill (1965) to allow a causal relationship to be inferred from observed associations, and by using additional criteria including:

- ▶ **Absence of negative consequences:** The presence of unintended negative consequences can have a major impact on the adoption or expansion of interventions. For example, fear that needle and syringe programmes might be seen as condoning drug use or that it may lead to security problems or violent behaviour or attacks.
- ▶ **Feasibility of implementation and expansion:** Is it feasible to implement programmes in prisons in diverse settings, including resource-poor settings, and in prisons of various types and security classifications, including in prisons for women?
- ▶ **Acceptability to the target of the intervention:** Do prisoners and staff accept the programmes and what conditions facilitate acceptance?
- ▶ **Unanticipated benefits:** Does the introduction of such programmes lead to other unintended and welcome benefits?

While the reliability of research conclusions without support from randomized clinical trials is often questioned, the difficulty of conducting such trials to evaluate public health interventions such as needle and syringe programmes in prisons should not be underestimated (e.g. Drucker et al, 1998). Generally, for a number of reasons, very few randomized clinical trials to evaluate HIV interventions in prisons have been undertaken.

1. EVIDENCE REGARDING INJECTING DRUG USE AND RESULTING TRANSMISSION OF HIV AND OTHER BLOOD-BORNE INFECTIONS IN PRISONS

1.1 Background

Illegal drugs are available in prisons despite the sustained efforts of prison systems to prevent drug use by prisoners by undertaking efforts to prevent the entry of drugs into prisons, by tightly controlling distribution of prescription medications, and enforcing criminal prohibitions on illegal drug possession and use among prisoners.

Many prisoners come to prison with established drug habits (Calzavara et al., 2003). Hiller et al. (1999) report that in the United States, 68% of all new admissions test positive for an illegal drug in urine screening, and similar findings have been reported across Europe (EMCDDA, 2005), North America, and Australia (Shewan, Stöver & Dolan, 2005). In other parts of the world, the situation is less clear because of the lack of systematic research (Dunn et al., 2000; Ohaeri, 2000), but in many countries, drug use among prisoners is common. In fact, many prisoners are in prison in the first place because of offences related to drugs (UNAIDS, 1997). These may be crimes related to drug production, possession, trafficking or use, or crimes committed to acquire resources to purchase drugs. Many prison systems have seen large increases in their population (and consequent overcrowding) attributable in large measure to a policy of actively pursuing and imprisoning those dealing with and consuming illegal substances (Stöver et al., 2001).

In particular for injecting drug users, imprisonment is a common event, with studies from a large number of countries reporting that between 56% and 90% of injecting drug users had been imprisoned at some stage (Ball et al. 1995; Normand et al. 1995; Millson, 1991; Wood et al., 2004; Beyrer et al., 2003). Multiple episodes of imprisonment are more common for IDU prisoners than for other prisoners (Gore et al., 1995). In a number of studies (Dolan, 2000), the percentage of prisoners with a history of injecting drug use before incarceration ranged from 11% in one study in England (Maden et al, 1992) to 64% in studies in New South Wales, Australia (Dolan et al., 1999).

People who used drugs prior to imprisonment often find a way to continue using on the inside, although prevalence and frequency rates for most – but not all (Plourde and Brochu, 2002; Swann & James, 1998) – prisoners decline with imprisonment (Shewan et al., 1994). Some people discontinue using drugs while in prison, while other prisoners start using drugs, often as a means to release tensions and to cope with being in an overcrowded and often violent environment (Taylor et al., 1995; Hughes & Huby, 2000). Plourde & Brochu (2002) found that drug use was significantly higher in maximum- (52%) and medium-security (35%) than in minimum-security institutions in Canada (19%). Cocaine use diminished considerably, while a significant number of prisoners who had not previously used heroin tried it in prison. This is consistent with findings of other studies revealing the popularity of heroin in prison (Swann & James, 1998).

Bullock (2003) found that the main reason provided by prisoners for their reduced levels of drug use in prisons was the relative lack of availability in prison (mentioned by 61% of those reporting reduced use), followed by attempts to stay off drugs (14%) and “get fit” (Kevin, 2000), not being able to afford drugs (13%), and concerns about punishment (6%: Bullock, 2003).

1.2 Injecting drug use in prison

Injecting drug use in prison is of particular concern with regard to transmission of HIV and other blood borne infections such as hepatitis B and C. This is because those who inject drugs in prisons often share needles and syringes and other injecting equipment (see *infra*, section 1.2.2), which is a very efficient way of transmitting HIV.

Studies may underestimate the prevalence of injecting drug use in prisons because of the many methodological, logistical, and ethical challenges of undertaking a study of prisoners’ high-risk behaviours. Injecting drugs is a highly clandestine activity (Hughes, 2000a), and many prisoners decline to participate in studies because they claim not to have

engaged in any high-risk behaviours (Health Canada, 2004, with reference to Pearson, 1995). This can result in low generalizability and underreporting of risk behaviours affecting statistics in prisons worldwide. As well, prisoners who do participate can be reluctant to give information regarding risk behaviours and, in particular, injecting drug use (Health Canada, 2004). Prisoners are afraid of reprisal for admitting illegal behaviours (Rutter, 2001, with reference to Dolan, Wodak & Penny, 1995).

In addition, caution must be exercised when comparing the prevalence of injecting and injecting risk behaviour between prisons in different countries, since studies have used different methodologies and indicators. However, despite these challenges, there is ample evidence that injecting drug use is widespread in prisons and represents a serious risk of HIV (and/or HCV) transmission.

As shown in **Table 1**, a large number of studies from countries around the world report high levels of injecting drug use, including among female prisoners (DiCenso, Dias & Gahagan, 2003; Elwood Martin et al., 2005). Studies also show that

- ▶ the extent and pattern of injecting and needle sharing vary among prisons
- ▶ many people who inject before imprisonment reduce or stop injecting when they enter prison, but many resume injecting upon release
- ▶ some people start injecting in prison; and
- ▶ those who inject in prison will usually inject less frequently than outside, but are much more likely to share injecting equipment than are drug injectors in the community (Shewan et al., 1994)
- ▶ those who inject in prison are sharing injection equipment with a population – fellow prisoners – that often has a high rate of HIV and HCV infections.

Most of these studies were undertaken in developed countries, but there are data from a number of developing countries and countries in transition (see **Table 1** and Dolan et al., 2007). For example, Rowhani-Rahbar, Tabatabee-Yazdi & Panahi (2004) report that about 10% of Iranian prisoners are believed to inject drugs and more than 95% of them are reported to share needles. Injecting in prison is a serious problem in prisons in countries in Eastern Europe and Central Asia (Russia: Frost & Tchertkov, 2002; Drobniowski et al., 2005; Ukraine: Zhivago, 2005; Armenia: Weilandt, Eckert & Stöver, 2005; Tajikistan: Godinho, 2005), and there are

also reports of injecting drug use in prisons in Latin America (e.g., Mexico: Cravioto P et al., 2003) and Africa (e.g., Rapid Situation Assessment Mauritius, 2005; Adjei et al., 2006).

1.2.1 Starting to inject in prison

Studies in prisons in many countries have reported that a relatively high percentage (13 to 23%) of people who inject in prison have started injecting in prison, for example in Irish prisons (Allright et al., 2000: 21% of injectors); Scottish prisons (Gore SM et al., 1995; Gore SM et al., 1997; Bird AG et al., 1997: 19% in one prison); Finland (Korte et al., 1998: 21.7%), Thailand (Thaisri et al., 2003: of 351 injectors, 15.9% initiated injecting while incarcerated), Russia (Frost & Tchertkov, 2002: 13.5%), Canada (Calzavara et al., 1997: 23%; Ford et al., 2000: 16%), and Australia (Dolan & Wodak, 1999). In other studies, the proportion of IDUs who started injecting in prison was somewhat lower (Bird AG et al., 1997: 4%; Bird AG et al., 1995: 6%; Power et al., 1992: 8% of a sample of male injectors in Scottish prisons). An overview prepared for the pre-expansion European Union reported that between 0.4% and 21% of injecting drug users started injecting in prison (EMCDDA, 2002).

Gill, Noone, and Heptonstall (1995) have suggested that the observation that a large number of prisoners begin injecting in prison should be interpreted with caution, saying that “if men who are at risk of becoming injecting drug users spend a substantial part of their young adult life in prison, the rate at which young men in prison become drug injectors may be no different from that for men of the same age outside prison.”

1.2.2 Using non-sterile injecting equipment in prison

Studies show that those who inject in prison are typically much more likely to share injecting equipment than are injecting drug users in the community (see **Table 1**), with most studies reporting sharing rates of between 60% and 90%. Because it is more difficult to smuggle needles and syringes into prisons than it is to smuggle drugs into them, needles and syringes are often in short supply. Often, only a handful of needles and syringes will circulate among a large population of prisoners who inject drugs. As a result, 15 to 20 people may inject using the same equipment (Correctional Service Canada, 1994; Small et al., 2005; Taylor & Goldberg, 1996). Sometimes, the equipment is home-made, and needle substitutes are fashioned out of hardened plastic and ball-point pens, often causing damage to veins, scarring, and

severe infections (Small et al., 2005; Mahon, 1996; Hughes, 2003; Turnbull, Stimson & Stillwell, 1994; Taylor & Goldberg, 1996; Bijl & Frost, 2000). In addition to the serious risk of infection, drug injectors in prison are at more risk of health complications, including scarring and bruising, abscesses and thrombosis from using extremely poor quality injecting equipment (Morrison, Elliott & Gruer, 1997).

In many cases, the successful record of risk reduction in the community contrasts with fairly stable reports of high risk using non-sterile injecting equipment in prison. In one study, over 15% of participants who reported injecting and sharing when last in prison also reported that was the first time they had ever shared injecting equipment (Crofts et al., 1995). Studies in Ireland, Scotland, and Australia reported that between 9.7% and 45.7% of those who were injecting prior to imprisonment reported having shared injecting equipment, but between 52.1% and 76% of those who were injecting in prison (Shewan et al., 1994; Allright et al., 2000; Kevin, 2000). The rate of sharing injecting equipment was particularly high among female prisoners (EMCDDA, 2002). Shewan et al. (1994) identified a number of factors significantly associated with current sharing of injecting equipment in prison: having injected a wider range of drugs in prison; frequency of Temgesic® (buprenorphine) use; and being prescribed methadone in the community, then having that prescription discontinued on entry into prison. Only one Canadian study found that the rates of injecting with non-sterile injecting equipment in prison were the same as pre-incarceration (Calzavara et al., 2003).

Use of non-sterile injecting equipment in prisons resembles that occurring in shooting galleries in that numerous strangers share syringes randomly in prison (Dolan, Wodak, Hall, Kaplan, 1998; Small, 2005). Generally, only friends or sexual partners share syringes in the community (Dolan et al, 1996a). The sharing that occurs in shooting galleries and in prisons is much more risky than other kinds of sharing and the difference is more pronounced when HIV prevalence is low (Allard, 1990). In prisons, interpersonal relationships and the possession of exchangeable resources determine access to scarce syringes. The scarcity of syringes results in patterns of sharing amongst large numbers of persons. In a study by Dolan et al. (1996a), 51 respondents outside prison shared syringes with 144 others; in prison, 60 respondents reported sharing with a total of 1,144 IDUs. Such continual reuse of scarce syringes poses serious health hazards (Small et al., 2005).

A small number of qualitative studies has examined HIV risk associated with injecting and sharing in prisons (Taylor & Goldberg, 1996; Hughes, 2003; Small et al., 2005). They report that used syringes may circulate for long periods and are used by many prisoners, and that sharing injecting equipment is difficult to avoid for prisoners who do inject because syringes are so scarce. Accessing syringes normally entails some form of payment unless a prisoner shares a close social relationship, like a friendship, with the owner. Ownership of injecting equipment can confer privileged position inside prison. It enables owners to levy a charge to others for the use of injecting equipment or trade drugs for the loan of injecting equipment. Some prisoners in the studies suggested that a prisoner may not disclose the fact that they are HIV positive, for fear that they would not be able to gain access to a syringe in future.

1.2.3 Determinants of injecting drug use in prison

A number of studies have found that drug use in prison is, at least partly, the product of a prison regime in which drugs are used in an attempt to combat boredom and isolation (Calzavara et al., 1997; Hughes & Huby, 2000). "It is important to recognise that the role of drugs in people's lives provides a meaningful social and self-identity inside prison, alleviates boredom, and fills the void that the absence of constructive regimes leaves" (Hughes, 2003, with reference to Her Majesty's Chief Inspector of Prisons for England and Wales, 1993; Hughes and Huby, 2000).

Studies have shown that motives for drug use prior to and during incarceration are quite different. Plourde & Brochu (2002) found that the majority of prisoners who had used drugs while in prison had used them to relax (62%), while prior to incarceration they had used drugs primarily to forget their problems (38%) and to have fun (31%). Calzavara et al. (1997) found that the top reasons for using drugs in the 12 months prior to incarceration were: "it makes me feel good", "because I'm addicted", and "a way to escape reality". In contrast, the top reasons for using drugs in the past 12 months of incarceration were: "it makes me feel good", "it makes the time pass easier", and "it helps me deal with feelings of boredom".

Furthermore, one study found that independent correlates of drug injecting while incarcerated were injection of heroin (OR=6.4) or other opiates (OR=7.9) and not injected with used needles (OR=0.20) outside in the year prior to incarceration; and ever being incarcerated in a federal prison (OR=5.3) (Calzavara

et al., 2003). A few other studies have suggested that prison-based drug use has more to do with the nature of the population and their pre-prison behaviour than the prison environment (Kevin, 2000; Thomas & Cage, 1975). Generally, there is agreement about the need for further research towards understanding why some prisoners maintain or even increase drug use in prison (see, e.g., Shewan, Stöver & Dolan, 2005). Health Canada (2004) points out that research studies undertaken to date often “lack more in-depth details regarding the motivations behind risk behaviours, which could aid in more effective planning and implementation of preventive measures” and suggests that future research should aim to identify the motivations of the prison population in engaging in high-risk conduct rather than elucidating specific behaviours and factors. This approach could help develop more tailored and effective prevention and intervention initiatives”.

1.2.4 Injecting upon release

As mentioned above, many drug using prisoners, including injectors, stop using drugs upon incarceration and are physically and behaviourally healthier while in prison than when in the community. But there is evidence of a high number of relapses (or taking up the old using patterns) and overdoses after this period of abstinence. In one study, relapse to drug injecting during the week following release from prison was reported by 41% of study participants, in 82% of cases on the very day of release (Van Haastrecht, Anneke & Van Den Hoek, 1998). This is consistent with the results of other studies reporting that a majority of prisoners who stop injecting in prison said they “definitely intend to” or “will probably” inject when released (Strang et al., 1998; Shewan et al., 2001).

1.3 Evidence of HIV and HCV transmission

A large number of studies from countries in many regions of the world have reported HIV and/or HCV and/or hepatitis B virus (HBV) seroconversion within prisons or shown that a history of imprisonment is associated with prevalent and incident HIV and/or HCV and/or HBV infection among IDUs.

With regard to HIV infection, it was significantly associated with a history of imprisonment in a number of countries in Western and Southern Europe (including among female prisoners: Estebanez et al., 2000), but also in Russia, Canada, Brazil, Iran, and Thailand. Using non-sterile injecting equipment in prison was found to be the most important independent deter-

minant of HIV infection in a number of studies (see **Table 2** and the chapter on “Evidence of the risk of HIV and HCV transmission in prisons” in the comprehensive paper on *Effectiveness of Interventions to Address HIV in Prisons* for more details).

The strongest evidence of extensive HIV transmission through injecting drug use in prison has emerged from a number of documented outbreaks in Australia (Dolan & Wodak, 1999), Lithuania (MacDonald, 2005), Russian Federation (Bobrik et al., 2005) and Scotland (Taylor et al., 1995). In the first documented outbreak, at least thirteen prisoners became infected at Glenochil prison in Scotland by using non-sterile injecting equipment (Taylor & Goldberg, 1996; Yirrell et al., 1997). A follow up study 12 months after the outbreak estimated that up to 20 prisoners had become infected (Gore et al., 1995).

In Lithuania, using non-sterile injecting equipment resulted in one of the largest documented HIV outbreaks in a prison. In May-June 2002, the Correctional Affairs Department and the Lithuanian AIDS Centre identified 207 HIV-positive prisoners at Alytus correctional facility. The survey was repeated in July 2002 and a further 77 HIV-positive prisoners were identified, of whom 44 had been found to be HIV-negative in May 2002. In total, 299 new HIV-positive cases were identified (MacDonald, 2005). A similar outbreak was also documented in a correctional colony in Tatarstan, Russian Federation, where 260 prisoners became HIV-infected in 2001 (Bobrik, 2005). Outbreaks of HIV have also been reported from other countries, but little information is available about these outbreaks (Dolan et al., 2007).

Finally, *HCV infection* by sharing of injecting equipment in prison has been reported in studies undertaken in Australia (Haber et al., 1999; O’Sullivan et al., 2003) and in Germany (Keppler, Nolte, Stöver, 1996; Keppler & Stöver, 1999).

2. EVIDENCE REGARDING PRISON-BASED NEEDLE AND SYRINGE PROGRAMMES

2.1 Background

Due to the prevalence of injecting drug use in prisons in many countries and the resulting risk of HIV and HCV transmission, providing sterile needles and syringes to prisoners has been widely recommended.

As early as 1993, in its *Guidelines on HIV Infection and AIDS in Prisons*, WHO recommended that “in countries where clean syringes and needles are made available to injecting drug users in the community, consideration should be given to providing clean injection equipment during detention and on release” (WHO, 1993). The same recommendation was made by UNAIDS (1997a; 1997b) and many other national and international bodies, including the Australian Medical Association (Editor, 1996) and the Ontario Medical Association (2004). The *International Guidelines on HIV/AIDS and Human Rights* also state that prison authorities should provide prisoners with the means for HIV prevention, including “clean injection equipment” (Office of the United Nations High Commissioner for Human Rights and UNAIDS, 2006, at 29e).

The rationale for establishing NSPs in prisons where injecting drug use takes place is even stronger than in communities (Rutter et al., 2001). Although injecting drug use in prison is usually less frequent than in the community, each episode involves more risk due to the scarcity of sterile injecting equipment and the higher prevalence of sharing of injecting equipment. Furthermore, the rapid turnover of prison populations means that there are more changes in injecting partners than in community settings; and also results in considerable interaction between prisoner- and community-based injecting drug user populations (Dolan, Rutter & Wodak, 2003).

The first prison NSP was established in Switzerland in 1992. Since then, NSPs have been introduced (or are about to be introduced), in over 50 prisons in 12 countries in Western and Eastern Europe and in Central Asia (see **Table 3**). In some countries, only a few prisons have NSPs, but in Kyrgyzstan and Spain NSPs have been rapidly scaled up and operate in a large number of prisons.

NSPs were first introduced in small prisons in Switzerland, but have since been implemented in other countries in prisons for men and for women;

in small, medium, and large institutions; in prisons of all security classifications; in civilian and military prisons; in different forms of custody (remand and sentenced, open and closed); and in institutions that house prisoners in individual cells as well as in those that house prisoners in barracks. Significantly, after having been introduced in well resourced prison systems in Western Europe, programmes have since been established in systems with very limited financial resources. Several models for the distribution of sterile injecting equipment have been used, including automatic dispensing machines; hand-to-hand distribution by prison physicians, other prison health care staff or drug counsellors, or by external community health workers; and distribution by prisoners trained as peer outreach workers. A brief overview of the history of prison NSPs can be found in the comprehensive paper on *Effectiveness of Interventions to Address HIV in Prisons* (available at <http://www.who.int/hiv/idu/>).

2.2 Evidence of effectiveness of NSPs in community settings

In many countries NSPs have become an integral part of a pragmatic public health response to reduce the risk of HIV transmission among injecting drug users and ultimately, to the general public. Up to 2007, some 60 countries have implemented legal and/or government sponsored NSPs in community settings.

Extensive studies have found NSPs to be effective in reducing HIV spread (General Accounting Office, 1993; Normand, Vlahov & Moses, 1995; Office of Technology Assessment of the US Congress, 1995; Institute of Medicine of the National Academy of Science, 2001; National Academy of Sciences, 2006). WHO has concluded that “measured against any objective standards, the evidence to support the effectiveness of NSPs in substantially reducing HIV must be regarded as overwhelming” (WHO, 2004, at 28).

2.3 Evidence of the effectiveness of NSPs in prison

Systematic evaluations of the effects of NSPs on HIV-related risk behaviours and of their overall effectiveness in prisons have been undertaken in at least 10 projects in Germany, Spain and Switzerland.

These evaluations were either of one or two years in duration, collecting data through a variety of means, and followed generally accepted scientific standards. Limitations include relatively small sample sizes, relatively short follow-up timeframes, inconsistent methodologies for assessing seroprevalence and seroincidence, and absence of comparison groups (Public Health Agency of Canada, 2006). Summaries of the most relevant results are provided in **Table 4**.

In addition, a study on the feasibility of NSPs in prisons was conducted in New South Wales, Australia, in 1995 (Rutter et al., 1995). Finally, while there are no published evaluations of NSPs in Eastern Europe and Central Asia, a number of published and unpublished reports, papers and presentations provide information about these NSPs and their effects.

The following questions guided the review and analysis of published and unpublished data on the effectiveness of prison-based NSPs.

- (1) Do prison NSPs lead to decreased risk behaviours among IDUs and are these changes in behaviour associated with lower rates of infection among IDUs in prison?
- (2) Do prison NSPs have additional and worthwhile benefits?
- (3) Is there any evidence of any major, unintended negative consequences?

2.3.1 Reduction in the use of non-sterile injecting equipment and of resulting blood-borne infections

With one exception (Heinemann & Gross, 2001), all available evaluations have shown that using non-sterile injecting equipment either ceased after implementation of the NSP (see, e.g., Nelles et al., 1998; Stark et al., 2005) or significantly dropped (see, e.g., Nelles, Fuhrer & Vincenz, 1999; Menoyo, Zulaica & Parras, 2000; Stöver, 2000) (see also **Table 4**). Injecting drug users in Moldovan prisons with NSPs also reported few incidents of sharing injecting equipment (Pintilei, 2005).

Due to the findings of ethical committees that comparison of different groups with and without access to NSPs would be unethical, the studies undertaken could not provide conclusive evidence of the impact of the NSPs on the incidence of blood-borne viral infections. However, no new cases of HIV were reported in any evaluation. In five of the six prisons in which blood tests were performed for HIV or hepatitis infection, no seroconversion was observed (summarized

in Stöver & Nelles, 2003), and self-reports in other prisons also indicated no new cases of infection. In another prison in which the incidence of HIV, HBV, and HCV was determined through repeated testing, no HIV and HBV seroconversions were observed, but four HCV seroconversions (Stark et al., 2005), one of which had definitely occurred in prison and was attributed to frontloading¹ (Stark et al., 2005).

Only in the evaluation of a NSP in the open prison of Hamburg-Vierlande, Germany, prisoners interviewed as part of a qualitative investigation reported only a small reduction in the use of non-sterile injecting equipment. Sharing continued because of insufficient supply with needles and syringes, mainly due to frequent break downs of the distribution machines, but also because the location of the machines did not allow for anonymous access, provision of the dummies that allowed for usage of the machines was inadequate, and because syringes of a particular size that was in high demand were not provided. However, the medical research team that conducted a quantitative investigation of prisoners' injecting behaviours reported more positive findings, including a much reduced rate of using non-sterile injecting equipment. In addition, no seroconversions were observed after the introduction of the NSP, while retrospective analysis of data recorded before the introduction of the NSP detected five hepatitis B and two hepatitis C seroconversions in the study group that must have happened in prison (Heinemann & Gross, 2001).

Overall, the observed reduction in the use of non-sterile injecting equipment is significant.

2.3.2 Additional benefits

There is evidence of ancillary health and social benefits associated with the implementation of NSPs.

Reduction in overdose incidents and deaths

A significant reduction of overdose incidents and deaths was reported in the first needle exchange pilot projects in Germany (Jacob & Stöver, 2000a; Jacob & Stöver, 2000b). Lines et al. (2004; 2005) also reported similar findings in the Hindelbank prison, Switzerland, which had averaged between one and three overdose deaths a year prior to the introduction of the NSP. In contrast, in the nine years after the NSP began, only one prisoner died of an overdose. Two reasons why NSPs have resulted in a decrease in overdose incidents and deaths have been offered:

¹ Dividing up drug doses between two or more injecting drug users involved in syringe sharing or sharing of spoons for drug preparation.

- ▶ Providing each injecting drug user with his/her own injecting equipment enables the individual to consume a smaller amount of drug with each injection. In the past, when a syringe was shared among many prisoners, a person who injected drugs would only have limited access to it and would be more likely to inject large doses on those rare occasions when he/she was in possession of the syringe.
- ▶ The implementation of NSPs and the adoption of a harm reduction philosophy within the prison fundamentally changes the way that prison health and social work staff are able to engage in counselling with prisoners. Honest discussions about risk behaviours and overdose risk can take place in an atmosphere in which prisoners do not have to fear sanctions for admitting drug use (Lines et al., 2004; 2005).

Increase in referral to drug treatment programmes

Evaluations of NSPs in Germany and Spain showed that they facilitated greater prisoner contact with drug treatment programmes, with referrals to drug treatment increasing (Stöver, 2000; Menoyo, Zulaica, Parras, 2000).

Other benefits

A number of evaluations noted other benefits, such as reduction in abscesses, a reduction in stress, improved relationships between prisoners and staff, and increased awareness of infection transmission and risk behaviours (Menoyo, Zulaica, Parras, 2000; and the summary in Lines et al., 2004; Lines et al., 2005).

There are also reports of increased staff safety in prisons with NSPs, due to the fact that accidental injuries from hidden injecting equipment during cell searches have decreased (Jürgens, 1996; Lines et al., 2004). Rihs-Middel (cited in Rutter et al., 1995) suggested that the decrease in the risk of injury is due to the fact that prisoners are permitted to store injecting equipment in a particular area of their cell and therefore do not hide it, reducing the risk of needle-stick injury during searches. Meyenberg et al. (1999) found that prison staff believed that the introduction of NSPs made injecting equipment easier to control.

With one exception, evaluation studies report no problems with the safe disposal of used syringes, and exchange rates of injecting equipment have been high, reaching 98.9 and 98.3 percent respectively in two German prisons (Meyenberg et al., 1999). Only

in the Hamburg-Vierlande prison there were reports of syringes not being disposed of properly. This was explained, at least in part, by two facts: prisoners felt that they would suffer negative consequences if they kept their syringe in the designated location; and access to sterile injecting equipment was limited (Heinemann & Gross, 2001).

2.3.3 Absence of unintended negative consequences

No serious unintended negative consequences were reported.

Syringes not used as weapons

Among the most important findings from the evaluation studies is that there was no reported instance where prisoners have used syringes as weapons against other prisoners or staff. Since the first NSP started in 1992, there have been no reports of syringes ever having been used as weapons in any prison with an operating NSP. The only report of a syringe ever being used as a weapon is from a prison in New South Wales, Australia, which did not have a NSP. In that case, a prison guard was stabbed with a blood-filled syringe by a HIV-positive prisoner, and subsequently seroconverted and died (Rutter et al., 2001; Jones, 1991).

NSPs do not lead to increased drug use or injecting

As outlined in **table 4**, evaluations of existing NSPs have found that the availability of sterile injecting equipment does not result in an increased number of injecting drug users, an increase in overall drug use or an increase in the amount of drugs in prisons (Jacob & Stöver, 2000a; Menoyo, Zulaica, Parras, 2000; Nelles, Fuhrer, Vincenz, 1999; Stark et al., 2005; Stöver, 2000; Stöver & Nelles, 2003). Evaluations of two programmes actually found that reported levels of drug use or injecting decreased (Nelles, Dobler-Mikola, Kaufmann, 1997).

Despite this evidence, there continues to be concern from some individuals that providing needles and syringes in prisons could lead to increased injecting drug use. Because this claim has sometimes been used to oppose implementation of NSPs, the findings of some inconclusive studies are discussed here in more detail.

In one study (Stark et al., 2005), two individuals who had previously only inhaled heroin reported injecting drug use on single occasions. It could not be ruled out that the availability of sterile injecting equipment may have facilitated initiation of injecting drug use

for these two individuals, but the researchers concluded that it is more likely that this finding reflects the natural incidence of injecting drug use among inhalation heroin users in settings where peers frequently inject (Stark et al., 2005, with reference to Allright et al., 2000; Gore et al., 1995).

In a letter to the *British Medical Journal*, Langkamp (2000) claimed that the evaluation of one NSP (in the Hamburg-Vierlande prison) found that many prisoners who had stopped using drugs started using them again, and that many prisoners went from inhaling drugs back to injecting drug use “while sharing needles regularly”. An analysis of the reports by the sociological and medical research teams (Gross, 1998; Heinemann & Gross, 2001) does show that a higher percentage of prisoners reported injecting at Hamburg-Vierlande than at the closed institutions from which they were transferred to Vierlande. Furthermore, some prisoners reported that they were tempted to recommence injecting drug use, though they had previously switched to other forms of drug use as a result of their fear of HIV and HCV transmission, in the absence of an NSP. However, the evaluation of the NSP does not make a link between increased drug use and injecting in Hamburg-Vierlande (an open prison, compared to the closed institutions from which prisoners were transferred) and the existence of the NSP at the prison. In addition, the evaluation report, while stating that prisoners’ reports (that they could be tempted to go back to injecting drug use because of the NSP) need to be taken seriously, stresses that these reports need to “be interpreted with great caution since a change in the consumption behaviour can very easily be attributed to the syringe distribution machines so that the responsibility and the ‘fault’ can be given to others than the prisoners themselves” (Gross, 1998, translation from German original).

Further, a switch from inhaling or smoking to injecting drug use has also been noted in studies undertaken in prisons without NSPs and attributed to the low availability of heroin that encouraged the switch from smoking to injecting drug use, a more efficient mode of consumption (Long, 2004).

The fear that the introduction of NSPs in prisons could tempt some prisoners to return to or continue injecting drug use in prisons was also expressed in one of the reviews of prison NSPs, not based on any of the experiences with existing NSPs, but on a small study undertaken in a prison in Canada. In that study, one of 11 prisoners who reported having stopped injecting as a result of being arrested or imprisoned specifically stated that he stopped injecting due to his inability

to get sterile injecting equipment and his concern over HIV transmission (Thomas, 2005). According to Thomas, “this appears to suggest that the introduction of sterile needles in prison could lead a small number of injecting drug users who had given up injecting because of the lack of sterile needles to return to using drugs intravenously.” Thomas suggested that future evaluations of NSPs in prison should include ethnographic data collection for the assessment of these types of potential behavioural effects.

Smyth (2000) also speculated that, “[a]lthough there is no evidence that provision of needle exchange encourages individuals to start injecting in the community, implementation of such a service could cause many more of these established injectors to opt to continue injecting while in prison.” He expressed a concern that NSPs in prisons could increase the incidence of HCV if more injecting drug users decide to continue injecting due to the presence of an NSP, and if some of them share injecting equipment occasionally, despite the presence of the NSP. He urged that a better understanding of the factors that mediate the observed reduction of injecting in this setting [prisons] is needed” and suggested that research evaluations of NSPs in prison should measure the proportion of prison entrants with injecting drug use histories who continue to inject before the introduction of the NSP and then re-measure that proportion after the introduction of the NSP. Smyth conceded that “there should still be substantial health gain for the wide population of injecting drug users from the provision of the NSP in the prison” if the proportion of injecting drug users who continue injecting “only increases marginally” (Smyth, 2006).

In conclusion, while evaluations of NSPs should continue to monitor the impact of NSPs on drug use and injecting in prisons, the evidence thus far shows that NSPs do not lead to increased drug use and injecting. The few reported instances in which a small number of prisoners may have switched to injecting drug use could be attributable to other factors. Even if they were related to the easier access to injecting equipment, they would not substantially impact the potential health benefits documented in the evaluations of NSPs.

2.3.4 Other findings

Adequate access to NSPs and need for confidentiality and trust

Ensuring that all prisoners have easy and confidential access to NSPs and develop trust that they can access injecting equipment when they need it and without having to fear any negative consequences

is a key factor in ensuring their success. Evaluations have shown that prisoners are reluctant to use the NSP if they fear that accessing injecting equipment may result in negative consequences, either because they could be seen using a dispensing machine (Heinemann & Gross, 2001) or because they could only access the NSP through health care or other staff (Stöver, 2000). Technical failures of dispensing machines, leading to limited access to injecting equipment, were also noted (Stöver, 2000).

In one prison in which equipment was distributed through counselling staff and prisoners receiving opioid substitution therapy were excluded from the NSP, needles and syringes remained a commodity for trade in the prison. There was also reluctance to access the NSP due to the lack of anonymity and a fear that counsellors' knowledge of participants' drug consumption could affect parole (Meyenberg et al., 1997; Jacob & Stöver, 1997). In at least one prison, sharing of injecting equipment continued because syringes of a particular size which were in high demand were not available, highlighting that the injecting equipment provided needs to meet the prisoners' demand (Heinemann & Gross, 2001).

If prisoners have limited access to the programme, are not provided the right type of syringes, or lack trust in the programme, benefits for staff will also be reduced, as some prisoners will continue to hide needles and syringes, thus increasing the risk of needlestick injuries for staff (Heinemann & Gross, 2001).

The extent to which easy access, confidentiality and trust are important has been best demonstrated in Moldova, where only a small number of prisoners accessed the NSP when it was located within the health care section of the prison. It was only when prisoners could obtain injecting equipment from fellow prisoners, trained to provide harm reduction services, that the number of equipment distributed increased significantly i.e. 98.4 percent of prisoners reported easy access to injecting equipment (Pintilei, 2005; Lines et al., 2004; Canadian HIV/AIDS Legal Network, 2006). This suggests that in many prisons, distribution by prison nurses or physicians or even by non-governmental organizations or health professionals who come to the prison for this purpose will not be the best option, as many prisoners would not access the programme. In these prisons, distribution through peers has led to much greater access, without any unintended negative consequences (Pintilei, 2005; Wolfe, 2005; Lines et al., 2004; Lines et al., 2005).

Finally, distribution, rather than one-for-one exchange, guarantees greater access to injecting equipment, particularly for those prisoners who are reluctant to access the NSP themselves and prefer to have injecting equipment delivered by trusted peers, and where opening hours are limited.

Acceptance of NSPs by staff and prisoners

Experience has shown that prior to the implementation of NSPs, prison staff have to be convinced to accept or at least tolerate them. Nevertheless, once in place, acceptance increases and is generally high among staff, as well as among prisoners who use drugs and those who do not (Nelles & Fuhrer, 1995; Nelles et al., 1998; Meyenberg et al., 1999).

The one exception was the Hamburg-Vierlande prison, where staff attitudes towards the NSP did not improve. The evaluators concluded that the NSP should not be extended to all prisons until staff had a chance to actively participate in the development of a model that responds to the needs and reality of each prison (Heinemann & Gross, 2001). Staff attitudes towards the NSP were least positive in those prisons in which prisoners experienced problems accessing syringes and/or did not trust that they could obtain them without suffering negative consequences, leading to the continued illegal trade of syringes and, generally, to reduced benefits of the NSP (Heinemann & Gross, 2001).

2.4 Conclusions and recommendations

There is evidence that NSPs are feasible in a wide range of prison settings.

Overall, the review of the evidence demonstrates that prison NSPs are feasible in a wide range of prison settings: in men's and women's prisons, prisons of all security levels, small and large prisons, and in prisons in which prisoners live in units of individual cells and in barracks-style facilities. It also demonstrates that NSPs can be successfully implemented in countries in which prison systems are relatively well resourced, as well as in countries in which prisons operate with significantly less funding and infrastructural support, such as in Eastern Europe (Moldova, Belarus, and Ukraine) and Central Asia (Kyrgyzstan).

Prison-based NSPs appear to be effective in reducing needle sharing and resulting HIV infection.

There is strong evidence that the provision of sterile injecting equipment is readily accepted by injecting drug users in prisons and may contribute to a signifi-

cant reduction of syringe sharing over time. Based on the data available and extrapolating from the vast literature on community-based programmes, prison NSPs also appear to be effective in reducing resulting HIV infections.

Prison-based NSPs have additional and worthwhile benefits.

In particular, there are reports that NSPs in prison

- ▶ lead to reduced overdose risk and a decrease in abscesses
- ▶ facilitate referral to drug dependence treatment programmes (where available) and lead to an increase in the number of prisoners accessing such programmes.

There is no convincing evidence of any major, unintended negative consequences.

There is no evidence to suggest that prison-based NSPs have serious, unintended negative consequences. In particular,

- ▶ NSPs do not appear to lead to increased drug use or injecting;
- ▶ Injecting equipment are not used as weapons;
- ▶ NSPs do not appear to undermine abstinence-based programmes, as drugs have remained prohibited within prisons where NSPs are in place. Security staff remain responsible for locating and confiscating illegal drugs. However, it is recognized that if and when drugs find their way into the prison and are used by prisoners, the priority must be to prevent the transmission of HIV and HCV via unsafe injecting practices. Therefore, while drugs themselves remain illegal, injecting equipment that is part of the official NSP is not.

In order to be successful, prisoners need to have easy, confidential access to NSPs, and prisoners and staff should receive information and education about the programmes and be involved in their design and implementation.

The review also showed that there are a number of key determinants of the success of prison-based NSPs:

- ▶ easy and confidential access by prisoners to NSPs
- ▶ support by prison staff and prisoners, emphasizing the importance of information and education of both staff and prisoners about the programme and its expected benefits for prisoners, staff, and the public
- ▶ developing a mechanism for safe disposal of syringes, and

- ▶ involvement of staff and prisoners in the design and development of the programmes.

Additional research about prison-based NSPs would be beneficial if it leads to reducing gaps in evidence.

The review has shown that there are areas in which future evaluation studies could reduce gaps in research. Most importantly, NSPs in prisons in countries outside Western Europe have not been scientifically evaluated. Moldova has been collecting various data and is undertaking prevalence studies (Pintilei, 2005), but none of the programmes implemented outside Western Europe collected data before the programmes began or has attempted to more systematically gather research data. Gathering additional data would be important to inform the prison systems in Eastern Europe and Central Asia in which NSPs are increasingly being introduced.

For additional studies, Rutter et al. (1995) recommend using a two-year evaluation using multi-method strategies including: quantitative and qualitative interviews of prisoners and staff; testing prisoners for blood borne viral infections and drug use; and review of prison records for assaults and/or drug seizures. According to Thomas (2005), the “key tasks” in evaluating future pilot prison-based NSPs “is to collect reliable information on a wide-range of relevant indicators before, during and after implementation of the program, analyse any changes that can be attributed to the needle exchange program, compare the ‘positive’ and ‘negative’ effects, and make a determination as to whether the positive effects (e.g., a reduction in the amount of needle sharing and disease transmission, etc.) outweigh the negative effects (e.g., prisoners being introduced to, returning to, or increasing injecting because of the availability of needles, etc.).”

In the end, as the United States National Academy of Sciences’ Institute of Medicine stated in the context of its analysis of the evidence on NSPs in the community, it has to be recognized that “the improbability of being able to carry out the definitive study ... does not necessarily preclude the possibility of making confident scientific judgments.” Citing Bradford Hill, the Institute continued saying that “incomplete” scientific evidence “does not confer upon us a freedom to ignore the knowledge we already have, or to postpone the action that it appears to demand” (Normand, Vlahov & Moses, 1995, cited in WHO, 2004; National Academy of Sciences, 2006).

To reject prison-based NSPs, based on limitations of the design of the studies undertaken thus far, would ignore both the preponderance and pattern of the evi-

dence and be “both poor scientific judgment and bad public health policy” (WHO, 2004) Or, in the words of WHO Europe (2005): “The relatively little experience available appears to show that, where risks are great, such as in countries with high prevalence rates of HIV and hepatitis, carefully introducing a syringe- and needle-exchange programme would be justifiable based on the experience already available ... When prison authorities have any evidence that injecting is occurring, they should consider an exchange scheme, regardless of the current prevalence of HIV infection”.

It is therefore recommended that:

1. Prison authorities in countries experiencing or threatened by an epidemic of HIV infections among IDUs should introduce needle and syringe programmes urgently and expand implementation to scale as soon as possible.

The overall success of the evaluated prison-based NSPs and the other available data reviewed for this report present a compelling case that prison-based NSPs are feasible, and suggest that they reduce sharing of injecting equipment and the resulting spread of HIV infections. This suggests that similar programmes may be beneficial in any prison with a problem of injecting drug use and associated sharing of injecting equipment.

The higher the prevalence of injecting drug use and associated risk behaviour is in prison, the more urgent introduction of prison-based NSPs becomes.

Monitoring and evaluation is an important component of any programme. While pilot projects of prison-based NSPs may be important in allowing the introduction of these programmes and to overcome objections against such programmes, they should not delay the expansion of the programmes, particularly where there already is evidence of high levels of injecting in prisons.

2. Additional research about prison-based NSPs should be undertaken to address remaining knowledge gaps.

This review has demonstrated significant gaps in research. In particular, more research in resource-poor systems outside Western Europe could allow for more rapid expansion of NSPs in these systems. Research in other systems should be designed to address research gaps rather than replicate existing studies. Evaluation of pilot programmes may be justified if: (1) the evaluation takes place in settings that are sufficiently different from settings in which evaluations have already been undertaken; or (2) it addresses research gaps.

3. EVIDENCE REGARDING BLEACH AND DECONTAMINATION STRATEGIES

3.1 Background

Programmes providing bleach or other disinfectants for sterilizing needles and syringes to reduce HIV transmission among injecting drug users in the community were first introduced in San Francisco, United States, in 1986 (Normand, Vlahov, Moses, 1995). Such programmes have received support particularly in situations where opposition to NSPs has been strongest, including in prisons in most countries (Rutter et al., 2001).

The number of prison systems that make disinfectants – mainly in the form of bleach - available to prisoners has continued to grow, but already in 1991, 16 of 52 prison systems surveyed made them available, including in Africa and Central America (Harding & Schaller, 1992). In surveys undertaken in Europe, the proportion of prison systems that make bleach available rose from 28 percent in 1992 to 50 percent in 1997 (European Network on HIV/AIDS and Hepatitis Prevention in Prisons, 1997). Today, bleach or other disinfectants are available in many other prison systems, including in Australia, Canada, Indonesia, Iran, and some systems in Eastern Europe and Central Asia (Lines, 2002; Dolan, 1999; Canadian HIV/AIDS Legal Network, 2006).

3.2 Evidence of effectiveness of programmes providing bleach in community settings

WHO (2004, at 28) has concluded that the “evidence supporting the effectiveness of bleach in decontamination of injecting equipment and other forms of disinfection is weak.” WHO pointed out that the efficacy of bleach as a disinfectant for inactivating HIV has been shown in numerous laboratory studies; that higher concentrations of bleach, although not always necessary, are more effective; and that contact time with bleach and the presence of other matter, such as clotted blood in syringes, are also important factors influencing efficacy (WHO, 2004 at 9). However, notwithstanding the strength of the laboratory data, field studies have cast “considerable doubt on the likelihood that these measures could ever be effective in operational conditions” (WHO, 2004 at 28). They concluded that disinfection of needles with bleach appeared to offer no protection, or at best little protection, against HIV infection (Chaisson et al., 1987; Vlahov et al., 1991; Titus et al., 1994; Vlahov et al., 1994).

Moreover, two studies assessed the effect of bleach on hepatitis C virus (HCV) prevalence and neither found a significant effect of bleach on HCV seroconversion (Kapadia et al., 2002; Hagan et al., 2001). At best, one of the studies (Kapadia et al., 2002) suggests a small (and probably insignificant) reduction of HCV infection.

3.3 Evidence from studies undertaken in prisons

Only a small number of studies have evaluated programmes providing bleach or other disinfectants in prison, and even fewer have focused on the health effects of such programmes.

3.3.1 Reduction of risk behaviours and of infections

The first two studies to allow the independent monitoring of a prison bleach-distribution programme were undertaken in Australia. They found that most prisoners could obtain bleach and that virtually all prisoners who were using non-sterile injecting equipment reported cleaning the syringes with bleach (Dolan et al., 1994; Dolan et al., 1996b; Dolan, Wodak, Hall, 1998; Dolan, Wodak, Hall, 1999). The studies also found that there was a significant improvement in access to bleach between the first and second study. Other Australian studies also showed that, when bleach is made available, a significant proportion of injecting drug users in prison clean syringes with bleach, but rates in some prisons were significantly lower (Rutter et al, 2001).

An evaluation of harm reduction measures in the Canadian federal prison system also reported relatively easy access to bleach, though in a few prisons access was not discreet (Correctional Service of Canada, 1999). In contrast, in a small qualitative study designed to examine the health risks experienced by male prisoners who inject drugs in British Columbia, Canada, prisoners claimed that the supply and quality of bleach in prisons was inconsistent, and that bleach is not always kept in an appropriate, accessible location (Small, 2005).

While studies show that a significant number of prisoners will clean syringes with bleach if it is accessible, studies also highlight that conditions in prisons make it even more unlikely than in the community that injecting equipment will be effectively decontaminated with bleach. The research

team that conducted the evaluation in the Canadian federal prison system (Correctional Service of Canada, 1999) stated that it had “no confidence that the distribution of bleach alone will effectively reduce transmission of infection from Hepatitis or HIV”. It concluded that “because of the clandestine and furtive nature under which injection drug users operate in prison settings; of the primitive and make shift equipment used to inject drugs; and, of the tendency of injection drug users to “cut corners” when their cravings overcome their judgment, there is no guarantee that the use of bleach alone will effectively reduce transmission of infection from HIV or Hepatitis C.”

This is consistent with the findings of the other studies that examined prisoners’ use of bleach, which reported that only a small number of prisoners reported adopting recommended syringe cleaning guidelines (Dolan & Wodak, 1998); bleaching of equipment in prisons “does not occur consistently, and most likely bleaching is performed too quickly when it is done” (Small, 2005); and that, while most prisoners claimed always to clean used equipment, “because prisoners can be accosted at any moment by prison officers, injecting and cleaning is a hurried affair” (Taylor & Goldberg, 1996).

3.3.2 Safety and security

There are no reports of any serious safety or security problems related to bleach programmes in prisons. The only evaluation that examined whether there were any unintended consequences related to the distribution of bleach kits in prison reported that both prisoners and staff said that bleach had become a ‘fact of life’ in prisons. Interviews with staff indicated that, with a few exceptions, staff concerns in terms of safety have abated (Correctional Service of Canada, 1999).

3.4 Conclusions and recommendations

Disinfection and decontamination schemes in the community outside prisons are not supported by evidence of effectiveness. In prisons, effectiveness may be reduced even further.

The type of syringes available in prisons may be more difficult to effectively disinfect with bleach, prisoners may have problems accessing bleach, and cleaning is a time consuming procedure and prisoners may be reticent to engage in any activity that increases the risk that prison staff will be alerted to their drug use. As WHO Europe has pointed out, “prisoners are highly unlikely to spend 45 minutes

shaking the syringes to clean them while waiting to inject in some hidden corner of the prison. Bleach can therefore create a false sense of security between prisoners sharing paraphernalia. The effectiveness of disinfection procedures ... depends greatly on the method used. Effectiveness varies and disinfection is now regarded as a second-line strategy to needle- and syringe-exchange programmes” (WHO Europe, 2005).

Distribution of bleach or other disinfectants is feasible in prisons and does not compromise security.

Disinfectants (mainly in the form of bleach) have been made available in a wide range of prison systems in different parts of the world. No reports of any serious safety or security problems related to bleach programmes could be found.

Because of their limited effectiveness, bleach programmes can only be regarded as a second-line strategy to NSPs. Therefore:

- Bleach programmes should be available in prisons where authorities continue to oppose the introduction of NSPs despite evidence of their effectiveness, and to complement NSPs. However, they cannot replace NSPs.
- Where bleach programmes are implemented, bleach should be made easily and discreetly accessible to prisoners in various locations in the prison, together with information and education about how to clean injecting equipment and information about the limited efficacy of bleach as a disinfectant for inactivating HIV and particularly HCV.
- Where bleach programmes exist in prisons, but not NSPs, public health practitioners should continue to advocate for the introduction of NSPs.

Table 1: Examples of studies that have examined injecting behaviour in prison

Location	Nr	% injected	% shared	Reference
Australia	2,482	36	60	Wodak 1989
Australia (NSW)	7 studies	31-74	70-94	Dolan & Wodak, 1999, with further references
Australia (SA)	50	52	60	Gaughwin, Douglas & Wodak 1991
Canada	4,285	11		Correctional Service Canada 1996
Canada	350	18.3		Ford et al. 2000
Canada	105 f	19		DiCenso, Dias, Gahagan 2003
Canada	102	21	86	Elwood Martin et al 2005
Canada	>1,200	27	80	Small et al., 2005
Canada	439 m, 158f	3.3	32	Calzavara et al., 2003
Canada	450	2.4	92	Dufour et al 1996
England	378	11.6	73	Edwards, Curtis, Sherrard, 1999
Europe (cross-sectional: France, Germany, Italy, Netherlands, Scotland, Sweden)	871	13		Rotily et al 2001b
European Union & Norway		0.2-34		EMCDD, 2005
Greece	544	24.1	92	Malliori et al 1998
Greece	861	20.2	83	Koulierakis et al 1999
Ireland	1178		70.5	Allright et al. 2000
Mauritius	100 m, 50 f, 50 youth (25 m, 25f)	10.8 of adults, 2.1 of youth		RSA Mauritius, 2005
Netherlands	497 injecting drug users	3	0	Van Haastrecht et al., 1998
Russian Federation	1,044	10	66	Frost & Tchertkov, 2002
Russian Federation	277	13		Dolan, Bijl & White, 2004
Scotland		15.9		Gore et al. 1995
Thailand	689	25	77.8	Thaisri et al. 2003
United States	281 m, 191 f	31% of injecting drug users with history of imprisonment had used illegal drugs in prison, and nearly half of these had injected in prison		Clarke et al. 2001

Table 2: Association of HIV, HCV, and HBV among injecting drug users with a history of imprisonment

Eastern Europe	Study	Finding
Russian Federation	Heimer et al., 2005	In this study of 826 injecting drug users in various cities in the Russian Federation, 44.8% reported ever having been to prison. Four health factors were correlated with imprisonment (HIV-positive; TB-positive, overdose and abscesses), while three were not (STIs, HBV and HCV).

Latin America	Study	Finding
Brazil	Varella et al., 1996	In this study of 82 male transvestites imprisoned in Sao Paulo, the factors associated with significant differences in positivity were the time spent in prison and the number of sexual partners during the previous year.
Brazil	Kallas et al., 1998	In this study of 780 prisoners in Sao Paulo, multivariate logistic regression analysis identified previous incarceration as an independent risk factor for HIV seropositive.
Brazil	Marins et al., 2000	In this study of prevalence and risk factors for HIV among 1059 prisoners, the number of previous incarcerations (1 compared to 0) (<i>OR</i> = 1.7, 95% <i>CI</i> 1.07–2.7) was an independent predictor of HIV.
Brazil	Guimaraes et., 2001	In this study of 779 prisoners of a prison in Sao Paolo, a time of current imprisonment longer than 130 months and previous incarceration at the same prison were associated with a positive anti-HCV serological test.
Brazil	Hacker et al., 2005	609 active/ex-injecting drug users were recruited from different communities, interviewed, and tested for HIV. Among male long-term injectors, “to have ever injected with anyone infected with HIV” (<i>Adj OR</i> = 3.91; 95% <i>CI</i> 1.09-14.06) and to have “ever been in prison” (<i>Adj OR</i> = 2.56; 95% <i>CI</i> 1.05-6.24) were found to be significantly associated with HIV infection.

North America	Study	Finding
Canada	Tyndall et al., 2003	In this study of injecting drug users in Vancouver, having been incarcerated in the last six months was independently associated with a markedly elevated incidence of HIV infection
Canada	Hagan, 2003	This external evaluation of the data in Tyndall et al. (2003) suggested that 21% of HIV infections among injecting drug users in Vancouver between 1996 and 2001 may have been attributable to infection during incarceration (Hagan, 2003).
Canada	Wood et al., 2005	Behaviours that can directly contribute to HIV infection (syringe borrowing and lending) were strongly and independently associated with reports of recent incarceration
Canada	Calzavara et al., 2005	Having a previous federal incarceration was found to be a risk factor significantly associated with HIV and HCV infection among adult prisoners in the Ontario provincial prison system.
United States	Fox et al., 2005	In this study of HCV infection among prisoners in the California state correctional system, independent correlates of HCV infection among both injecting drug user and non-injecting drug user prisoners included cumulative time of incarceration.

EVIDENCE REGARDING BLEACH AND DECONTAMINATION STRATEGIES

Pacific, South and South-East Asia	Study	Finding
Australia	Butler et al., 1997	Among prisoners entering the New South Wales correctional system, multivariate analysis identified previous imprisonment as a significant predictor for HCV infection
Australia	Butler et al., 1999	Multivariate analysis identified injecting while in prison as a major risk factor for HBV, and institutionalization as a factor for HCV.
Australia	Van Beek et al., 1998	A history of imprisonment was found to be an independent predictor of HCV seroconversion; HCV incidence was substantially higher among injecting drug users who had been imprisoned (60,8/100 person years) than those who had not (12,5/100 person years) .
Australia	Hellard, Hocking, Crofts (2004)	HCV-positive prisoners were more likely to have injected drugs (OR 29.9) and to have injected drugs in prison during their current incarceration (OR 3.0); injecting drugs whilst in prison during this incarceration was a risk factor for HCV.
Australia	Gates et al., 2004	A history of prior imprisonment was a risk factor associated with HCV infection.
Islamic Republic of Iran	Zamani et al., 2005	Among male injectors visiting treatment centres in Tehran, a history of shared injection inside prison (adjusted odds ration (OR, 12.37; 95% CI, 2.94-51.97) was the main factor associated with HIV-1 infection.
Thailand	Choopanya et al., 1991	Bangkok injecting drug users with a history of prison were about twice as likely to be HIV-positive as those who had never been jailed.
Thailand	Kitayaporn et al., 1998	Concluded that Bangkok injecting drug users continue to be at high risk for HIV infection related to use of non-sterile injecting equipment and incarceration.
Thailand	Vanichseni et al., 2001	In a cohort of injecting drug users in Bangkok, people who injected while incarcerated had a higher incidence of HIV infection (35.3 per 100 person years of observation) than those who had been incarcerated but had not injected (11.3 per 100) and those who had not been incarcerated (4.9 per 100). The authors concluded that the "great risk associated with incarceration warrants special attention. Although the risk associated with incarceration is not fully characterized, it is likely that a large proportion of this risk results from the use of non-sterile injecting equipment in settings where access to clean syringes and needles is severely limited."
Thailand	Beyrer et al., 2003	This study reaffirmed the association between incarceration and HIV infection among Thai male and female injecting drug users.

Western Europe	Study	Finding
England & Wales	Weild et al., 2000	Presence of anti-HCV was associated with injecting drug use inside prison and number of previous times in prison
France	Richardson et al., 1993	Imprisonment associated with HIV infection.
Germany	Stark & Muller, 1993; Muller et al., 1995; Stark et al., 1995a; 1995b; 1997	The use of non-sterile injecting equipment in prison was the most important independent determinant of HIV infection among a sample of injecting drug users in Berlin, and also an important determinant of HBV and HCV infection.

Greece	Malliori et al., 1998	The use of non-sterile injecting equipment in prison, and multiple imprisonments are the most important risk factors for HCV infection in injecting drug users.
Greece	Koulierakis et al., 2000	In this study among prisoners in 10 Greek prisons, logistic regression analysis suggested that total time in prison, previous drug conviction, being a convict (as opposed to on remand) and having multiple female sexual partners 1 year before incarceration were significant HIV risk behaviour correlates. For every year of imprisonment, the risk of injection in prison increased by about 17% [OR = 1.17 (95% CI: 1.07-1.27)]
Ireland	Allright et al., 2000	Time in prison over the past ten years and the use of non-sterile injecting equipment while in prison associated with HCV positivity. Concluded that "being in prison in Ireland may be an independent risk factor for contracting hepatitis C infection."
Italy	Babudieri et al., 2005	Frequency of imprisonment and tattoos were associated, respectively, with HIV and HCV positivity in a sample of prisoners from 8 Italian prisons.
Scotland	Davies et al., 1995	HIV infection was significantly associated with being imprisoned among a city-wide sample of injecting drug users in Edinburgh who had injected in the previous 6 months.
Scotland	Champion et al., 2004	Ever having injected drugs (relative risk= 13.0, 95% CI: 1.5, 114.3) and having shared needles/syringes in prison (relative risk= 9.0, 95% CI: 1.1, 71.7) were significantly associated with HCV seroconversion in prison.
Scotland	Seaman & Bird, 2001	No conclusive effect of incarceration on risk of HIV infection was found, but there was a suggestion that imprisonment might have been a significant relative risk factor for infection after risk behaviour among drug users in the community was reduced, due to introduction of NSPs.
Spain	Estebanez et al., 1990	Seropositivity increases with the number of times individuals are incarcerated.
Spain	Granados et al., 1990	Imprisonment associated with HIV infection.
Spain	Anon et al., 1995	HCV correlated with duration & number of imprisonments.
Spain	Pallas et al., 1999	Reincarceration and long-term injecting were the foremost risk factors for HBC-HCV and for HIV-HBV-HCV co-infection among injecting drug using prisoners.
Spain	Martin et al., 1998	Multiple incarceration histories and long-term imprisonment associated with higher risk of HIV infection.
Wales	McBride et al., 1994	HCV associated with history of imprisonment.
Multi-centre	Estebanez et al., 2000	In a multicentred, cross-sectional study undertaken in a population of female injecting drug users recruited from a variety of settings in Berlin,, London ,Madrid, Paris and Rome, factors independently associated with HIV prevalence in the regression analysis included previous imprisonment (OR = 1.4).

Table 3: Countries with needle and syringe programmes in prisons

Country	Start of programmes	Number of prisons with NSPs (as of 2006)
Switzerland	1992	7
Germany	1996	1 (6 NSPs were closed as a result of political decisions)
Spain	1997	38
Republic of Moldova	1999	7
Kyrgyz Republic	2002	11
Belarus	2003	1 (as of 2004)
Armenia	2004	3
Luxembourg	2005	1
Islamic Republic of Iran	2005	1 to 6 (five programmes to be opened in 2006)
Ukraine	2007	2 pilot projects expected to start in 2007
Scotland	2007	one 2-year pilot study approved for start in 2007
Portugal	2007-2008	implementation of NSPs by 2008

Table 4: Sample evaluations of needle and syringe programmes in prisons

Prison, Country	Incidence of HIV/HCV	Needle Sharing	Drug Use	Injecting
Am Hasenberge (D) (reported in Stöver & Nelles, 2003)	no data	strongly reduced	no increase	no increase
Basauri (E) (Menoyo, Zulaica, Parras, 2000)	no seroconversion	strongly reduced	no increase	no increase
Hannöversand (D) (reported in Stöver & Nelles, 2003)	no data	strongly reduced	no increase	no increase
Hindelbank (SUI) (Nelles, Dobler-Mikola, Kaufmann, 1997)	no seroconversion	strongly reduced	decrease	no increase
Berlin (Lehrter Strasse and Lichtenberg (Stark et al., 2005)		strongly reduced	no increase	no increase*
Lingen 1 (D) (Stöver, 2000; Jacob & Stöver, 2000a)	no seroconversion	strongly reduced	no increase	no increase
Realta (SUI) (Nelles, Fuhrer, Vincenz, 1999)	no seroconversion	single cases	decrease	no increase
Vechta (D) (Stöver, 2000; Jacob & Stöver, 2000a)	no seroconversion	strongly reduced	no increase	no increase
Vierlande (D) (Heinemann & Gross, 2001)	no seroconversion	little change or reduction	no increase	no increase

(adapted from Thomas, 2005; Stöver & Nelles, 2003; Rutter et al., 2001)

* 2 people who had previously only inhaled heroin reported injecting drug use on single occasions.

REFERENCES

- Adjei AA et al. (2006). Prevalence of human immunodeficiency virus, hepatitis B virus, hepatitis C virus and syphilis among prison inmates and officers at Nsawam and Accra, Ghana. *Journal of Medical Microbiology*, 55: 593–597.
- Allard R et al. (1990). A mathematical model to describe the risk of infection from sharing injecting equipment. *Journal on AIDS*, 3: 1010-1016.
- Allright S et al. (2000). Prevalence of antibodies to hepatitis B, hepatitis C, and HIV and risk factors in Irish prisoners: results of a national cross sectional survey. *British Medical Journal*, 321: 78-82.
- Anon C et al. (1995). The hepatitis C virus among the prison population in Valencia [article in Spanish]. *Rev Esp Enferm Dig*, 87(7): 505-508.
- Babudieri S et al. (2005). Correlates of HIV, HBV, and HCV infections in a prison inmate population: Results from a multicentre study in Italy. *Journal of Medical Virology*, 76 (3): 311-317.
- Ball A et al. (1995). Multi-centre Study on Drug Injecting and Risk of HIV Infection: a report prepared on behalf of the international collaborative group for World Health Organization Programme on Substance Abuse. Geneva: WHO.
- Beyrer C et al (2003). Drug use, increasing incarceration rates, and prison-associated HIV risks in Thailand. *AIDS and Behavior*, 7(2): 153-161.
- Bijl M, Frost L (2000). Establishing an HIV/AIDS prevention and health promotion program in Russian prisons. 11th International Conference on the Reduction of Drug Related Harm. Jersey, April 9-13.
- Bird AG et al. (1995). Anonymous HIV surveillance with risk factor elicitation at Scotland's largest prison, Barlinnie. *AIDS*, 9: 801-808.
- Bird AG et al. (1997). Harm reduction measures and injecting inside prison versus mandatory drugs testing: results of a cross sectional anonymous questionnaire survey. *British Medical Journal*, 315(7099): 21-24.
- Bobrik A et al. (2005). Prison health in Russia: the larger picture. *Journal of Public Health Policy*, 26: 30-59.
- Bradford Hill A (1965). The environment and disease: association or causation. *Proceedures of the Royal Society of Medicine*, 58: 295-300.
- Bullock T (2003). Changing levels of drug use before, during and after imprisonment. In: Ramsay M (ed). *Prisoners' drug use and treatment: seven research studies*. Home Office Research Study 267. London: Home Office Research, Development and Statistics Directorate.
- Butler T et al. (1997). Hepatitis B and C in New South Wales prisons: prevalence and risk factors. *Medical Journal of Australia*, 166: 127.
- Butler T et al. (1999). Seroprevalence of markers for hepatitis B, C and G in male and female prisoners - NSW, 1996. *Australian and New Zealand Journal of Public Health*, 23(4), 377-384.
- Calzavara L et al. (1997). *Understanding HIV-Related Risk Behaviour in Prisons: The Inmates' Perspective*. Toronto: HIV Social, Behavioural and Epidemiological Studies Unit, Faculty of Medicine, University of Toronto.
- Calzavara LM et al. (2003). Prior opiate injection and incarceration history predict injection drug use among inmates. *Addiction*, 98: 1257-1265.
- Canadian HIV/AIDS Legal Network (2006). HIV/AIDS in Prisons in Central and Eastern Europe and the former Soviet Union. Bleach and other disinfectants (Info sheet 5). Montreal: The Network.
- Chaisson RE et al. (1987). HIV, bleach, and needle-sharing. *Lancet*, 1(8547): 1430.
- Champion J et al. (2004). Incidence of hepatitis C virus infection and associated risk factors among Scottish prison inmates: a cohort study. *American Journal of Epidemiology*, 159: 514-519.
- Choopanya K et al. (1991) Risk factors and HIV seropositivity among injecting drug users in Bangkok. *AIDS*, 5: 1509-1513.
- Clarke JG et al. (2001). Active and former injection drug users report of HIV risk behaviors during periods of incarceration. *Subst Abuse*, 22(4): 209-216.
- Correctional Service Canada (1994a). *HIV/AIDS in Prisons: Final Report of the Expert Committee on AIDS and Prisons*. Ottawa: Minister of Supply and Services Canada.
- Correctional Service Canada (1996). *1995 National Inmate Survey: Final Report*. Ottawa: CSC (Correctional Research and Development), No SR-02.

- Correctional Service Canada (1999). *Evaluation of HIV/AIDS Harm Reduction Measures in the Correctional Service of Canada*. Ottawa: CSC.
- Cravioto P et al. (2003). Patterns of heroin consumption in a jail on the northern Mexican border: barriers to treatment access. *Salud Publica de Mexico*, 45: 181-190.
- Crofts N et al. (1995). Spread of bloodborne viruses among Australian prison entrants. *British Medical Journal*, 310: 285-288.
- Davies A et al. (1995). HIV and injecting drug users in Edinburgh: Prevalence and correlates. *Journal of Acquired Immune Deficiency Syndrome Human-Retroviral*, 8: 399-405.
- DiCenso A, Dias G, Gahagan J (2003). *Unlocking Our Futures: A National Study on Women, Prisons, HIV, and Hepatitis C*. Toronto: PASAN.
- Dolan K et al. (1994). *Bleach Availability and Risk Behaviours in New South Wales*. Technical Report No 22. Sydney: NDARC.
- Dolan K, Wodak A, Penny R (1995). AIDS behind bars: preventing HIV spread among incarcerated drug injectors. *AIDS*, 9: 825-832.
- Dolan K, et al. (1996a). HIV Risk Behaviour of IDUs before, during and after Imprisonment in New South Wales. *Addiction Research*, 4(2): 151-160.
- Dolan K et al. (1996b). *Bleach Easier to Obtain But Inmates Still at Risk of Infection in New South Wales Prisons*. Technical Report. Sydney: NDARC.
- Dolan K, Wodak AD (1998). A bleach program for inmates in NSW: an HIV prevention strategy. *Aust N Z J Public Health*, 22(7): 838-840.
- Dolan K, Wodak A, Hall W (1998). Methadone maintenance treatment reduces heroin injection in NSW prisons. *Drug and Alcohol Review*, 17(2): 153-158.
- Dolan K, A Wodak, W Hall, E Kaplan (1998). A mathematical model of HIV transmission in NSW prisons. *Drug & Alcohol Dependence*, 50: 197-202.
- Dolan K (1999). The epidemiology of hepatitis C infection in prison populations. National Drug and Alcohol Research Centre, UNSW.
- Dolan K, Wodak A (1999). HIV transmission in a prison system in an Australian State. *Medical Journal of Australia*, 171(1): 14-17.
- Dolan K, Wodak A, Hall W (1999). HIV risk behavior and prevention in prison: a bleach program for inmates in NSW. *Drug and Alcohol Review*, 18: 139-143.
- Dolan K (2000). The epidemiology of hepatitis C infection in prison populations. In: *Hepatitis C: Informing Australia's National Response*. Canberra: Commonwealth Department of Health and Aged Care.
- Dolan K, Rutter S, Wodak A (2003). Prison-based syringe exchange programmes: a review of international research and development. *Addiction*, 98, 153-158.
- Dolan K et al. (2004). Review of injection drug users and HIV infection in prisons in developing and transitional countries. UN Reference Group on HIV/AIDS Prevention and Care among IDUs in Developing and Transitional Countries.
- Dolan J, Kite B, Aceijas C, Stimson GV (2007). HIV in prison in low-income and middle-income countries. *Lancet Infectious Diseases*, 7 : 32-43.
- Dolan K, Bijl M, White B (2004). HIV education in a Siberian prison colony for drug dependent males. *International Journal of Equity in Health*, 3: 7.
- Drobniewski FA et al. (2005). Tuberculosis, HIV seroprevalence and intravenous drug abuse in prisoners. *Eur Respir J*, 26(2): 298-304.
- Drucker E et al. (1998). Measuring harm reduction: the effects of needle and syringe exchange programmes and methadone maintenance on the ecology of HIV. (Review). *AIDS*, 12 (Suppl. A): S217-230.
- Dufour A Alary M, Poulin C et al. (1996). Prevalence and risk behaviours for HIV infection among inmates of a provincial prison in Quebec City. *AIDS*, 10:1009-1015.
- Dunn J, Laranjeira R, Marins J (2000). HIV, drug use, crime, and the penal system: competing priorities in a developing country – the case of Brazil. In: Shewan D & Davies JB (eds). *Drug Use and Prisons: An International Perspective*. Chur: Harwood.
- Editor (1996). Australian Medical Association calls for needle exchange programs for prisoners. *Canadian HIV/AIDS Policy & Law Newsletter*, 2(4): 25.
- Edwards A, Curtis S, Sherrard J (1999). Survey of risk behaviour and HIV prevalence in an English prison. *Int J STD AIDS*, 10(7): 464-466.

- Elwood Martin R et al. (2005). Drug use and risk of bloodborne infections: A survey of female prisoners in British Columbia. *Canadian Journal of Public Health*, 96(2): 97-101.
- Estebanez PE et al. (1990). Jails and AIDS. Risk factors for HIV infection in the prisons of Madrid. *Gaceta sanitaria*, 4(18): 100-105.
- Estebanez PE et al. (2000) Women, drugs and HIV/AIDS: results of a multicentre European study. *International Journal of Epidemiology*, 29: 734-43.
- European Network on HIV/AIDS and Hepatitis Prevention in Prisons (1997). Final Report on the EU Project European Network on HIV/AIDS Prevention in Prisons. Bonn and Marseille: The Network.
- European Monitoring Centre on Drugs and Drug Addiction (EMCDDA)(2002). *2002 Annual Report on the State of the Drugs Problem in the European Union and Norway*. Luxembourg: Office for Official Publications of the European Community.
- European Monitoring Centre on Drugs and Drug Addiction (EMCDDA) (2005). *The State of the Drugs Problem in Europe. Annual Report 2005*. Luxembourg: Office for Official Publications of the European Community.
- Ford PM et al. (2000). HIV, hepatitis C and risk behaviour in a Canadian medium-security federal penitentiary. *Quarterly Journal of Medicine*, 93: 113-119.
- Fox et al. (2005). Hepatitis C virus infection among prisoners in the California state correctional system. *Clinical Infectious Diseases*, 41(2): 177-186.
- Frost L, Tchertkov V (2002). Prisoner risk taking in the Russian Federation. *AIDS Education and Prevention*, 14 (Suppl B): 7-23.
- Gates J et al. (2004). Risk factors for hepatitis C infection and perception of antibody status among male prison inmates in the hepatitis C incidence and transmission in prisons study cohort, Australia. *Journal of Urban Health*, 81(3): 448-452.
- Gaughwin MD, Douglas RM, Wodak AD (1991). Behind bars – risk behaviours for HIV transmission in prisons, a review. In: J Norberry, SA Gerull & MD Gaughwin (eds). *HIV/AIDS and Prisons* (conference proceedings). Canberra: Australian Institute of Criminology.
- General Accounting Office (1993). *Needle exchange programs: research suggests promise as an AIDS prevention strategy*. Washington DC: United States Government Printing Office.
- Gill O, Noone A, Heptonstall J (1995). Imprisonment, injecting drug use, and bloodborne viruses (editorial). *British Medical Journal*, 310: 275-276.
- Godinho J. (2005). Reversing the Tide: Priorities for HIV/AIDS Prevention in Central Asia. Washington: The World Bank.
- Gore SM et al. (1995). Drug injection and HIV prevalence in inmates of Glenochil Prison. *British Medical Journal*, 310: 293-296.
- Gore SM, Bird AG, Ross A (1995). Prison rites: Starting to inject inside. *British Medical Journal*, 311: 1135-1136.
- Gore SM et al. (1997). Anonymous HIV surveillance with risk factor elicitation at Perth (for men) and Cornton Vale (for women) prisons in Scotland. *International Journal on STDs and AIDS*, 8: 166-175.
- Granados et al. (1990). HIV seropositivity in Spanish prisons. Presented at the VIth International AIDS Conference, San Francisco. Abstract no Th.D.116.
- Gross U (1998). *Wissenschaftliche Begleitung und Beurteilung des Spritzenaustauschprogramms im Rahmen eines Modellversuchs der Justizbehoerde der Freien und Hansestadt Hamburg. Evaluationsbericht eines empirischen Forschungsprojekts*. Kriminologisches Forschungsinstitut Niedersachsen. [Evaluation of the prison needle exchange program in Hamburg]
- Guimaraes T et al. (2001). High prevalence of hepatitis C infection in a Brazilian prison: identification of risk factors for infection. *Brazilian Journal of Infectious Diseases*, 5(3): 111-118.
- Haber PS et al. (1999). Transmission of hepatitis C within Australian prisons. *Medical Journal of Australia*, 171: 31-33.
- Hacker MA et al. (2005). The role of “long-term” and “new” injectors in a declining HIV/AIDS epidemic in Rio de Janeiro, Brazil. *Subst Use Misuse*, 40(1): 99-123.
- Hagan H. (2003). The relevance of attributable risk measures to HIV prevention planning. *AIDS*, 17: 911-913.


- Harding TW, Schaller G (1992). *HIV/AIDS and Prisons: Updating and Policy Review. A Survey Covering 55 Prison Systems in 31 Countries*. Geneva: WHO Global Programme on AIDS.
- Health Canada – Public Health Agency of Canada (2004). Hepatitis C Virus Transmission in the Prison/ Inmate Population. *Canada Communicable Disease Report*, 30(16): 141-148.
- Heimer R et al. (2005). Imprisonment as risk for HIV in the Russian Federation: evidence for change. 16th International Conference on the Reduction of Drug Related Harm.
- Heinemann A, Gross U (2001). Prevention of blood-borne virus infections among drug users in an open prison by vending machines. *Sucht* 2001; 47(1): 57-65.
- Hellard ME, Hocking JS, Crofts N (2004). The prevalence and the risk behaviours associated with the transmission of hepatitis C virus in Australian correctional facilities. *Epidemiology Infect*, 132(3): 409-15.
- Her Majesty's Chief Inspector of Prisons for England and Wales (1993). *Doing Time or Using Time. Presented to Parliament by the Secretary of State for the Home Department by Command of Her Majesty, January 1993* (Cm 2128). London, Her Majesty's Stationery Office.
- Hiller ML, Knight K & Simpson D (1999). Prison-based substance abuse treatment, residential after-care and recidivism. *Addiction*, 94(6): 833-842.
- Hughes RA (2000a). Drug injectors and prison mandatory drug testing. *Howard Journal Of Criminal Justice*, 39(1): 1-13.
- Hughes RA (2000b). Lost opportunities? Prison needle and syringe exchange schemes. *Drugs: Education, Prevention and Policy*, 7(1): 75-86.
- Hughes RA, Huby M (2000). Life in prison: Perspectives of drug injectors. *Deviant Behavior*, 21(5): 451-479.
- Hughes RA (2003). Illicit drug and injecting equipment markets inside English prisons: a qualitative study. *Journal of Offender Rehabilitation*, 37(3/4): 47-64.
- Inciardi JA & Lockwood D (1993). Drug use in prison: Patterns, processes and implications for treatment. *Journal of Drug Issues*, 23(1), 119-130.
- Institute of Medicine of the National Academy of Science (2001). *No time to lose: getting more from HIV prevention*. Washington DC: National Academies Press.
- Jacob J, Stöver H (1997). Germany - needle exchange in prisons in Lower Saxony: a preliminary review. *Canadian HIV/AIDS Policy & Law Newsletter*, 3(2/3): 30-31.
- Jacob J, Stöver H (2000a). Drug use, drug control and drug services in German prisons: Contradictions, insufficiencies and innovative approaches. In: Shewan D, Davies J (eds). *Drug Use and Prisons: An International Perspective*. Amsterdam: Harwood Academic Publishers, 57-88.
- Jacob J, Stöver H (2000b). The transfer of harm-reduction strategies into prisons: needle exchange programmes in two German prisons. *International Journal of Drug Policy*, 11: 325-335.
- Jones PD (1991). HIV transmission by stabbing despite zidovudine prophylaxis (letter). *Lancet*, 338: 884.
- Jürgens R (1996). *HIV/AIDS in Prisons: Final Report*. Montréal: Canadian HIV/AIDS Legal Network and Canadian AIDS Society, at 52-66.
- Kallas EG et al (1998). HIV seroprevalence and risk factors in a Brazilian prison. *Braz J Infect Dis*, 2(4): 197-204.
- Kapadia F et al. (2002). Does bleach disinfection of syringes protect against hepatitis C infection among young adult injection drug users? *Epidemiology*, 13(6): 738-741.
- Keppler K, Nolte F, Stover H (1996). Transmission of infectious diseases in prison: results of a study in the prison for women in Vechta, Lower Saxony, Germany. Originally published in German in *Sucht*, 42(2): 98-107.
- Keppler K, Stöver H. (1999) Transmission of infectious diseases during imprisonment – results of a study and introduction of a model project for infection prevention in Lower Saxony. *Gesundheitswesen*, 61(4): 207-213. Summarized in English in *Canadian HIV/AIDS Policy & Law Newsletter* 1996; 2(2), 18-19.
- Kevin M (2000). *Addressing the Use of Drugs in Prison: A Survey of Prisoners in New South Wales*. Sydney: NSW Department of Corrective Services (Research Publication No. 44).

- Kitayaporn D et al (1998). Infection with HIV-1 subtypes B and E in injecting drug users screened for enrollment into a prospective cohort in Bangkok, Thailand. *Journal of Acquired Immune Deficiency Syndromes and Human Retrovirology*, 19: 289-295.
- Korte T, Pykalainen J, Seppala T (1998). Drug abuse of Finnish male prisoners in 1995. *Forensic Sci*, 97(2-3): 171-183.
- Korte T, Pykalainen J, Seppala T (1998). Drug abuse of Finnish male prisoners in 1995. *Forensic Sci*, 97(2-3): 171-183.
- Koulierakis G et al. (2000). HIV risk behaviour correlates among injecting drug users in Greek prisons. *Addiction*, 95(8):1207-16.
- Langkamp H (2000). Risks of syringe exchange programmes in prisons prevail. *British Medical Journal*, 321: 1406-1407.
- Lines R. (2002) *Action on HIV/AIDS in Prisons: Too Little, Too Late – A Report Card*. Montreal: Canadian HIV/AIDS Legal Network.
- Lines R et al. (2004). *Prison Needle Exchange: A Review of International Evidence and Experience*. Montreal: Canadian HIV/AIDS Legal Network.
- Lines R et al. (2005). Taking action to reduce injecting drug-related harms in prisons: The evidence of effectiveness of prison needle exchange in six countries. *International Journal of Prisoner Health* 1(1): 49-64.
- Long J, Allwright S, Begley C (2004). Prisoners' view of injecting drug use and harm reduction in Irish prisons. *International Journal of Drug Policy*, 15(2): 139-149.
- MacDonald M (2005). *A Study of Health Care Provision, Existing Drug Services and Strategies Operating in Prisons in Ten Countries from Central and Eastern Europe*. Finland: Heuni.
- Maden A, Swinton M & Gunn J (1993). A survey of pre-arrest drug use in sentenced prisoners. *British Journal of Addiction*, 87: 27-33.
- Mahon N (1996) New York inmates' HIV risk behaviors: the implications for prevention policy and programs. *American Journal of Public Health*, 86: 1211-1215.
- Malliori M et al. A survey of bloodborne viruses and associated risk behaviours in Greek prisons. *Addiction* 1998; 93(2): 243-251.
- Marins JR et al (2000). Seroprevalence and risk factors for HIV infection among incarcerated men in Sorocaba, Brazil. *AIDS and Behavior*, 4(1): 121-128.
- Martin V et al. (1998) Predictive factors of HIV-infection in injecting drug users upon incarceration. *European Journal of Epidemiology*, 14(4): 327-331.
- McBride AJ, Ali IM, Clee W (1994). Hepatitis C and injecting drug use in prisons. *British Medical Journal*, 309: 876.
- Menoyo C, Zulaica D, Parras F (2000). Needle exchange in prisons in Spain. *Canadian HIV/AIDS Policy & Law Review*, 5(4): 20-21.
- Meyenberg R, Stöver H, Jacob J, Pospeschill M (1999). *Infektionsprophylaxe im Niedersächsischen Justizvollzug*. Oldenburg: BIS-Verlag, 418 pp.
- Millson P (1991). Evaluation of a programme to prevent HIV transmission in injection drug users in Toronto. Toronto: Toronto Board of Health.
- Morrison A, Elliott L & Gruer L (1997). Injecting-related harm and treatment-seeking behaviour among injecting drug users. *Addiction*, 92(10): 1349-1352.
- Muller R et al. (1995). Imprisonment: A risk factor for HIV infection counteracting education and prevention programmes for intravenous drug users. *AIDS*, 9: 183-190.
- National Academy of Sciences (Committee on the Prevention of HIV Infection among Injecting Drug Users in High-Risk Countries) (2006). *Preventing HIV Infection among Injecting Drug Users in High Risk Countries: An Assessment of the Evidence*. Washington, DC: National Academies Press.
- Nelles J, Fuhrer A (1995). *Drug and HIV prevention at the Hindelbank penitentiary. Abridged report of the evaluation results of the pilot project*. Berne: Swiss Federal Office of Public Health.
- Nelles J, Dobler-Mikola A, Kaufmann B (1997). Provision of syringes and prescription of heroin in prison. The Swiss experience in the prisons of Hindelbank and Oberschöngrün. In: Nelles J, Fuhrer A (eds). *Harm Reduction in Prison*. Bern: Peter Lang, 239-262.

- Nelles J et al. (1998) Provision of syringes: the cutting edge of harm reduction in prison? *British Medical Journal*, 317(7153): 270-273.
- Nelles J, Fuhrer A, Vincenz I (1999). *Evaluation der HIV- und Hepatitis-Prophylaxe in der Kantonalen Anstalt Realta. Schlussbericht*. Berne: Universitäre Psychiatrische Dienste Bern.
- Nelles J, Fuhrer A, Hirsbrunner HP (1999). How does syringe distribution affect consumption of illegal drugs by prisoners? *Drug and Alcohol Review*, 18(2): 133-138.
- Normand J, Vlahov D, Moses LE, eds (1995). *Preventing HIV transmission: the role of sterile needles and bleach*. Washington DC: National Academy Press.
- Office of Technology Assessment of the US Congress (1995). *The effectiveness of AIDS prevention efforts*. Washington DC: US Government Printing Office.
- Office of the United Nations High Commissioner for Human Rights and UNAIDS (2006). *HIV/AIDS and Human Rights: International Guidelines* (2006 consolidated version). Geneva: UNAIDS (HR/PUB/06/9).
- Ohaeri JU (2000). Drug use and HIV/AIDS in sub-Saharan African Prisons. In: Shewan D & Davies JB (eds). *Drug Use and Prisons: An International Perspective*. Chur: Harwood.
- Ontario Medical Association (2004). Improving Our Health: Why is Canada lagging behind in establishing needle exchange programs in prisons? Toronto: The Association.
- O'Sullivan B et al. (2003). Hepatitis C transmission and HIV post-exposure prophylaxis after needle- and syringe-sharing in Australian prisons. *Medical Journal of Australia*, 178(11): 546-549.
- Pallas JR et al. (1999). Risk factors for mono-infections and coinfections with HIV, hepatitis B and hepatitis C viruses in northern Spanish prisoners. *Epidemiol Infect*, 123: 95-102.
- Pearson M (1995). Voluntary screening for hepatitis C in a Canadian federal penitentiary for men. *Canadian Communicable Disease Report*, 21(14): F4-F5.
- Pintilei L (2005). Harm reduction in prisons of Republic of Moldova. Presentation at "HIV/AIDS in Prisons in Ukraine – From Evidence to Action: Prevention and Care, Treatment, and Support." Kiev, 1-2 November.
- Plourde C, Brochu S (2002). Drugs in prison: a break in the pathway. *Substance Use Misuse* 2002; 37: 47-63.
- Power KG, Markova I, Rowlands A, McKee KJ, Anslow PJ, Kilfedder C (1992). Intravenous drug use and HIV transmission amongst inmates in Scottish prisons. *British Journal of Addiction*, 87: 35-45.
- Public Health Agency of Canada (2006). *Prison needle exchange: Review of the evidence*. Ottawa: The Agency, April 2006.
- Rapid Situation Assessment Mauritius (2005). Unpublished paper on file with author.
- Richardson C, Ancelle-Park R, Papaevangelou G (1993). Factors associated with HIV seropositivity in European injecting drug users. *AIDS*, 7: 1485-1491.
- Rotily M et al. (2001) Surveillance of HIV infection and related risk behaviour in European prisons. A multicentre pilot study. *Eur J Public Health*, 11(3): 243-250.
- Rowhani-Rahbar A, Tabatabaee-Yazdi A, Panahi M (2004). Prevalence of common blood-borne infections among imprisoned injection drug users in Mashhad, North-East Iran. *Archives of Iranian Medicine*, 7(3): 190-194.
- Rutter S et al. (1995). *Is Syringe Exchange Feasible in a Prison Setting? An Exploration of the Issues*. Technical Report No 25. Sydney: National Drug and Alcohol Research Centre, 1995.
- Rutter S et al. (2001). *Prison-Based Syringe Exchange Programs. A Review of International Research and Program Development* (NDARC Technical Report No. 112). Sydney: National Drug and Alcohol Research Centre, University of New South Wales.
- Seaman SR, Bird SM (2001). Proportional hazards model for interval-censored failure times and time-dependent covariates: application to hazard of HIV infection of injecting drug users in prison. *Stat Med*, 20(12): 1855-70.
- Shewan D, Gemmell M, Davies JB (1994). Behavioural change amongst drug injectors in Scottish prisons. *Soc Sci Med*, 39(11): 1585-1586.

- Shewan D et al. (2001). Injecting risk behaviour among recently released prisoners in Edinburgh (Scotland): The impact of in-prison and community drug treatment services. *Legal and Criminological Psychology*, 6: 19-28.
- Shewan D, Stöver H, Dolan K (2005). Injecting in prisons. In: Pates R, McBride A, Arnold K (eds). *Injecting Illicit Drugs*. London: Blackwell Publishing, 69-81.
- Small W et al. (2005). Incarceration, addiction and harm reduction: inmates experience injecting drugs in prison. *Substance Use & Misuse*, 40: 831-843.
- Smyth B (2000). Health effects of prisons (letter). *British Medical Journal*, 321: 1406.
- Stark K, Muller R (1993). HIV prevalence and risk behaviour in injecting drug users in Berlin. *Forensic Sci Int*, 62(1-2): 73-81.
- Stark K et al (1995a). Determinants of HIV infection and recent risk behaviour among injecting drug users in Berlin by site of recruitment. *Addiction*, 90(10): 1367-1375.
- Stark K et al. (1995b). Prevalence and determinants of anti-HCV seropositivity and of HCV genotype among intravenous drug users in Berlin. *Scandinavian Journal of Infectious Diseases*, 27(4) 331-337.
- Stark K et al. (1997). History of syringe sharing in prison and risk of hepatitis B virus, hepatitis C virus, and human immunodeficiency virus infection among injecting drug users in Berlin. *International Journal of Epidemiology*, 26(6): 1359-1366.
- Stark K et al. (2005). A syringe exchange programme in prison as prevention strategy against HIV infection and hepatitis B and C in Berlin, Germany. *Epidemiol Infect*, Dec 22; 1-6 [Epub ahead of print].
- Stöver H (2000). Evaluation of needle exchange pilot project shows positive results. *Canadian HIV/AIDS Policy & Law Newsletter*, 5(2/3): 60-64.
- Stöver H et al. (2001). *An overview study: assistance to drug users in European Union prisons*. Lisbon: European Monitoring Centre for Drugs and Drug Addiction.
- Stöver H, Nelles J (2003). 10 years of experience with needle and syringe exchange programmes in European prisons: A review of different evaluation studies. *International Journal of Drug Policy*, 14: 437-444.
- Strang J et al. (1998). HIV/AIDS risk behaviour among adult male prisoners (Research Findings No. 82). London: Home Office Research, Development and Statistics Directorate.
- Swann R, James P (1998). The effect of the prison environment upon inmate drug taking behaviour. *Howard Journal of Criminal Justice*, 37: 252-265.
- Taylor A et al. (1995). Outbreak of HIV infection in a Scottish prison. *British Medical Journal*, 310(6975): 289-292.
- Taylor A, Goldberg D (1996). Outbreak of HIV infection in a Scottish prison: why did it happen? *Canadian HIV/AIDS Policy & Law Newsletter*, 2(3): 13-14.
- Thaisri H et al (2003). HIV infection and risk factors among Bangkok prisoners, Thailand : a prospective cohort study. *BMC Infectious Diseases*, 3: 25.
- Thomas C & Cage R (1975). Correlates of prison drug use: an evaluation of two conceptual models. Paper presented at the Southern Conference on Corrections, Tallahassee, Florida.
- Thomas G (2005). *Assessing the need for prison-based needle exchange programs in Canada: a situational analysis*. Ottawa: Canadian Centre on Substance Abuse.
- Titus S et al. (1994). Bleach use and HIV seroconversion among New York City injecting drug users. *Journal of Acquired Immune Deficiency Syndromes*, 7(7): 700-704.
- Turnbull PJ, Stimson GV & Stillwell G (1994). *Drug Use in Prison*. Horsham: AIDS Education and Research Trust.
- Tyndall et al. (2003). Intensive injection cocaine use as the primary risk factor in the Vancouver HIV-1 epidemic. *AIDS*, 17: 887-893.
- UNAIDS (1997a). *Prisons and AIDS: UNAIDS point of view*. Geneva: UNAIDS.
- UNAIDS (1997b). *Prisons and AIDS: UNAIDS technical update*. Geneva: UNAIDS.
- United Nations Office on Drugs and Crime, WHO, UNAIDS (2006). *HIV/AIDS Prevention, Care, Treatment, and Support in Prison Settings. A Framework for an Effective National Response*. New York: United Nations.

- Van Beek I et al. (1998). Infection with HIV and hepatitis C virus among injecting drug users in a prevention setting: retrospective cohort study. *British Medical Journal*, 317: 433-437.
- van Haastrecht H, Bax JS, van den Hoek AR (1998) High rates of drug use, but low rates of HIV risk behaviours among injecting drug users during incarceration in Dutch prisons. *Addiction*, 93(9): 1417-25.
- Vanichseni S et al. (2001). Continued high HIV-1 incidence in a vaccine trial preparatory cohort of injection drug users in Bangkok, Thailand. *AIDS*, 15: 397-405.
- Varella D et al (1996). HIV infection among Brazilian transvestites in a prison. *AIDS Patient Care STDS*, 10(5): 299-302.
- Vlahov D et al. (1991). Prevalence of antibody to HIV-1 among entrants to US correctional facilities. *Journal of the American Medical Association*, 265: 1129-32.
- Vlahov D et al. (1994). Field effectiveness of needle disinfection among injecting drug users. *Journal of Acquired Immune Deficiency Syndromes*, 7(7): 760-766.
- Weilandt C, Eckert J & Stöver H (2005). Anonymous survey on infectious diseases and related risk behaviour among Armenian prisoners and on knowledge, attitudes and behaviour of Armenian prison staff towards infectious diseases and drugs. Bonn: WIAD, ENDIPP, ICRC.
- Weild AR et al. (2000). Prevalence of HIV, hepatitis B, and hepatitis C antibodies in prisoners in England and Wales: a national survey. *Communicable Disease and Public Health*, 3(2): 121-126.
- Wolfe D (2005). *Pointing the Way: Harm Reduction in Kyrgyz Republic*. Bishkek: Harm Reduction Association of Kyrgyzstan "Partners' network".
- Wood et al. (2004). Inability to access addiction treatment and risk of HIV infection among injection drug users. *Journal of Acquired Immune Deficiency Syndrome*, 36: 750-754.
- Wood E et al. (2005). Recent incarceration independently associated with syringe sharing by injection drug users. *Public Health Reports*, 120: 150-156.
- World Health Organization (1993). *WHO guidelines on HIV infection and AIDS in prisons*. Geneva: WHO (WHO/GPA/DIR/93.3).
- World Health Organization (2004). *Evidence for Action Technical Papers: Effectiveness of Sterile Needle and Syringe Programming in Reducing HIV/AIDS among Injecting Drug Users*. Geneva: WHO.
- World Health Organization Europe (2005). *Status paper on prisons, drugs and harm reduction*. Copenhagen, WHO Europe.
- Yirrell D et al. (1997). Molecular investigation into outbreak of HIV in a Scottish prison. *British Medical Journal*, 314: 1446.
- Zamani S et al. (2005). Prevalence of an factors associated with HIV-1 infection among drug users visiting treatment centres in Tehran, Iran. *AIDS*, 19(7): 709-716.
- Zhivago S (2005). HIV/AIDS epidemic situation in penitentiary system of Ukraine. Presentation at "HIV/AIDS in Prisons in Ukraine – From Evidence to Action: Prevention and Care, Treatment, and Support." Kiev, 1-2 November.



ISBN 978 92 4 159581 0



9 789241 595810